

**Universidade de Lisboa**  
**Faculdade de Farmácia**



# **Pancreatic cancer. New screening biomarkers and detection methods**

**Mariana Amaral Colaço de Andrade Gaspar**

**Mestrado Integrado em Ciências Farmacêuticas**

**2019**



**Universidade de Lisboa**  
**Faculdade de Farmácia**



# **Pancreatic cancer. New screening biomarkers and detection methods**

**Mariana Amaral Colaço de Andrade Gaspar**

**Monografia de Mestrado Integrado em Ciências Farmacêuticas**  
**apresentada à Universidade de Lisboa através da Faculdade de**  
**Farmácia**

**Orientadora: Professora Doutora Ana Cristina Ferreira da**  
**Conceição Ribeiro, Professora auxiliar FFUL**

**2019**



## Resumo

O cancro do pâncreas posiciona-se em quarto lugar como o de maior letalidade, com mais de 90% dos doentes diagnosticados nos estadios III e IV. A taxa de sobrevivência de cinco anos é de apenas 8,5%, o que pode ser explicado pela ausência de sintomas específicos e pela falta de uma estratégia de diagnóstico padronizada, de sensibilidade e precocidade acrescida. Pelo facto de o tumor primário necessitar de pelo menos cinco anos para obter capacidade metastática, o diagnóstico *early-stage* (como forma de melhorar a qualidade de vida e sobrevivência dos pacientes com cancro pancreático) é possível. Urge assim, uma atualização baseada na especificidade e sensibilidade, na atualização de biomarcadores, facilmente implementados na estratégia laboratorial e, economicamente concretizável.

**Palavras-chave:** biomarcador, cancro pancreático, diagnóstico early-stage

# Abstract

Pancreatic cancer ranks fourth as the most lethal with more than 90% of patients being diagnosed in stages III and IV. The five-year survival rate is only 8.5%, which can be explained by the absence of specific symptoms and by the lack of a standardized guideline for an early-stage diagnostic strategy with high sensitivity. Due to the fact that the primary tumor requires at least five years to obtain metastatic capacity, early-stage diagnosis (as a way to improve the life and survival chance of pancreatic cancer patients) is possible. However, there is an urge for an update in specificity and sensitivity and an update in biomarkers easily implemented in laboratory strategy and economically feasible.

**Keywords:** biomarker, pancreatic cancer, early-stage diagnosis

# Agradecimentos

A realização desta tese (e todo o mestrado integrado) não teria sido possível sem o apoio incondicional da minha família, orientadora e amigos.

À professora Doutora Ana Cristina Ribeiro, agradeço por toda a sua disponibilidade, dedicação, tempo, energia e por tudo o que me ensinou. Obrigada por todo o seu apoio e orientação durante estes meses. Não teria sido possível sem a professora.

À minha família, obrigada pela paciência, pelo apoio e pela compreensão, por me terem proporcionado esta possibilidade, por saber que posso sempre contar convosco.

# Table of Contents

<b>1</b>	<b>INTRODUCTION.....</b>	<b>1</b>
<b>1.1</b>	<b>Pancreatic Cancer .....</b>	<b>2</b>
<b>1.2</b>	<b>Epidemiology .....</b>	<b>5</b>
<b>1.3</b>	<b>Etiology .....</b>	<b>8</b>
1.3.1	Non-modifiable risk factors .....	8
1.3.2	Modifiable risk factors .....	10
<b>1.4</b>	<b>Current Diagnosis Methods.....</b>	<b>13</b>
1.4.1	Multidetector computed tomography (MDCT) .....	14
1.4.2	Magnetic Resonance Imaging (MRI) .....	14
1.4.3	Endoscopic Ultrasonography with fine needle aspiration (EUS).....	15
1.4.4	Positron Emission tomography (PET).....	15
1.4.5	Biopsy.....	15
1.4.6	Blood test.....	15
<b>1.5</b>	<b>Therapeutics.....</b>	<b>16</b>
1.5.1	Surgery.....	16
1.5.2	Chemotherapy .....	16
1.5.3	Radiation Therapy.....	16
<b>2</b>	<b>PANCREATIC CARCINOGENESIS.....</b>	<b>16</b>
<b>2.1</b>	<b>Pathology .....</b>	<b>16</b>
<b>2.2</b>	<b>Genetics.....</b>	<b>17</b>
<b>3</b>	<b>CURRENT STATE OF THE ART IN DIAGNOSTICS.....</b>	<b>18</b>
<b>3.1</b>	<b>Novelties in Pancreatic Biomarkers.....</b>	<b>18</b>
3.1.1	Traditional tumor biomarkers.....	19
3.1.1.1	Carbohydrate Antigen 19-9 (CA19-9).....	19
3.1.1.2	Other conventional biomarkers .....	23
3.1.2	Metabolites.....	23
3.1.3	Cell-free DNA .....	24



3.1.3.1	Somatic mutations .....	25
3.1.3.2	Epigenetic modulations.....	26
3.1.4	Autoantibodies .....	28
3.1.5	Cell-free noncoding RNA (ncRNA).....	29
3.1.5.1	microRNA (miRNA).....	30
3.1.5.2	Long non-coding RNA (lncRNA).....	32
3.1.6	Inflammatory factors and growth factors.....	34
3.1.7	Circulating tumor cells (CTCs) .....	36
3.1.8	Exosomes .....	36
3.1.8.1	Proteins .....	37
3.1.8.2	DNA with somatic mutations .....	37
3.1.8.3	miRNAs.....	38
3.1.9	Soluble stroma-related Biomarkers .....	38
<b>3.2</b>	<b>Glycosylation in Cancer .....</b>	<b>40</b>
3.2.1	Glycomic aberration vs. biomarkers .....	41
<b>3.3</b>	<b>Lectins as a tool .....</b>	<b>47</b>
3.3.1	Biomedical Application .....	48
3.3.2	Immobilized-Lectin Affinity Chromatography.....	48
3.3.3	Enzyme-linked Lectin Assay .....	49
3.3.4	Lectin Histochemistry .....	50
3.3.5	Lectin Blotting.....	51
3.3.6	Lectin Array.....	52
<b>4</b>	<b>CONCLUSIONS.....</b>	<b>53</b>
<b>5</b>	<b>CHALLENGES AND FUTURE DIRECTIONS.....</b>	<b>56</b>
	<b>BIBLIOGRAPHY .....</b>	<b>57</b>
	<b>ANNEXES.....</b>	<b>96</b>
<b>A1.</b>	<b>The evidence for cancer risk: summary matrix .....</b>	<b>96</b>



# 1 Introduction

Cancer is one of the biggest public health problems worldwide, as it is the major cause of mortality and morbidity (Mordente *et al.*, 2015). The lifetime risk of developing cancer is of almost 40% (National Cancer Institute, 2018).

Cancer is a term defining a wide variety of pathologies that can occur in almost any part of the body. Its common properties include local invasion and distant spread.

It develops when there is an accumulation of genetic damage in cells over time, due to the failing of the normal corrective processes (figure 1). The cells acquire abnormal qualities, namely rapid cell division, immortality and invasion of surrounding tissue (World Cancer Research Fund/American Institute for Cancer Research, 2018).




Figure 1. Hallmarks of cancer

(adapted from Hanahan *et al.*, 2011, with permission from Elsevier)

There are many tumor causing factors. Its composition, evolution, diagnosis and treatment also vary according to its subtype (La Thangue *et al.* 2011).

There is currently still an imperative need to develop a new, sensitive, specific and easy way to enable early diagnosis (Makawita *et al.*, 2010). Tumor biomarkers are a

very promising, non-invasive and quick way for early diagnosis, staging and monitoring disease progression and therapeutic response.

## 1.1 Pancreatic Cancer

The pancreas is a gland localized behind the stomach, containing two different parts: exocrine and endocrine. While the exocrine pancreas is responsible for the production of digestive enzymes, the endocrine pancreas produces hormones, like insulin and glucagon, being therefore included in the metabolism of glucose (World Cancer Research Fund, 2018).

Pancreatic cancer is not a single disease but a large number of neoplasms being divided into the two main pancreas functions: exocrine and endocrine.

The majority of the pancreatic cancers develop in the exocrine pancreas (about 95%) with most of them being localized in the pancreas head (World Cancer Report, 2014), namely adenocarcinoma (Vincent *et al.*, 2011, Ferlay *et al.*, 2012, Hidalgo *et al.*, 2013). The most aggressive pancreatic neoplasm is the ductal adenocarcinoma (Hruban *et al.*, 2007). Nevertheless, the early-detection of ductal adenocarcinoma improves the overall survival (Matsuno *et al.*, 2004). Concerning the endocrine tumors, they make up for less than 5% of cases (Vincent *et al.*, 2011, Ferlay *et al.*, 2012, Hidalgo *et al.*, 2013) and have a better five-year survival rate (65%) (Hruban *et al.*, 2007). The most common benign pancreatic neoplasm is the serous cystadenoma (Tseng *et al.*, 2005). Almost all patients with this type of tumor are cured (World Cancer Report, 2014).

Pancreatic cancer is the fourth cancer with more related deaths (Siegel *et al.* 2018). Because of the diagnosis difficulties, it is normally not diagnosed in time, having an extremely high mortality and poor prognosis (Bunger *et al.*, 2011). According to Brierly *et al.* (2017), more than 90% of patients are diagnosed only at stages III and IV.

According to the National Cancer Institute (2018), the number of deaths caused by pancreatic cancer has recently outweighed the ones caused by breast cancer (table 1).

Table 1. Comparison of breast and pancreatic cancer

(adapted from National Cancer Institute, 2018)

Breast Cancer			Pancreatic cancer		
<b>At a Glance</b>			<b>At a Glance</b>		
Estimated New Cases in 2018	266,120	<b>Percent Surviving 5 Years</b> <b>89.7%</b> 2008-2014	Estimated New Cases in 2018	55,440	<b>Percent Surviving 5 Years</b> <b>8.5%</b> 2008-2014
% of All New Cancer Cases	15.3%		% of All New Cancer Cases	3.2%	
Estimated Deaths in 2018	40,920		Estimated Deaths in 2018	44,330	
% of All Cancer Deaths	6.7%		% of All Cancer Deaths	7.3%	

Due to the fact that there is a lack of standardized international guideline (Zhang *et al.*, 2018a) regarding its diagnostic strategy, even for high-risk populations (Bunger *et al.*, 2011), nor specific symptoms (World Cancer Research Fund, 2018), pancreatic cancer is normally diagnosed in metastatic stage (Stathis *et al.*, 2010). Metastasis is the most prevailing motive of death in cancer patients (Leary *et al.*, 2010). It consists in the dissemination and growth of the neoplastic cells away from the primary tumor (Yachida *et al.*, 2010).

After diagnosis, very few individuals survive five years, namely 8,5% (National Cancer Institute, 2018). These poor rates have not been improving in the last 40 years (figure 2) (Zhang *et al.*, 2018a).



Figure 2. Five-year survival rate

(adapted from National Cancer Institute, 2018)

Almost 80% of the diagnosed patients have spread cancer, either in regional or metastatic phase (Costello *et al.*, 2012), having no opportunity to go through a curative treatment, as the surgical resection (which is the only curative option) is only possible in early stages (Yachida *et al.*, 2010). According to the National Cancer Institute (2018), only 9,7% of diagnosed patients with pancreatic cancer are at local stage (figure 3). This majority of patients do not have symptoms or radiologically visible manifestations either as the disease progresses to the metastatic phase (Ghaneh *et al.*, 2007, Hidalgo *et al.*, 2010) making the early detection, with the current available methods, still very difficult. Nevertheless, the survival rates improve, if the pancreatic cancer is diagnosed,

and treated, at an early stage, confined to the primary site (Herrerros-Villanueva, *et al.* 2016).




Figure 3. Percent of cases by stage of diagnosis

(adapted from National Cancer Institute, 2018)

In addition to that, cancer screening programs for other cancers, like breast, colorectal, cervical and lung cancer, have proven to reduce its mortality rates (Hakamaet *et al.*, 1986, Lynge *et al.*, 1989, Mandel *et al.*, 1999, Sigurdsson *et al.*, 1999, Tabár *et al.*, 2000, Nyströmet *et al.*, 2002, Atkin *et al.*, 2010, Aberle *et al.*, 2011, Schoenet *et al.*, 2012, Hur *et al.*, 2016), due to the fact that screening programs enable and increase the likelihood of an early-stage diagnosis. In early-stage, the tumor is less aggressive and still not metastatic making the chances of undergoing through curative measures majorly higher.

According to Yachida *et al.* (2010) and Poruk *et al.* (2013), in pancreatic cancer there are required at least five years, for the primary tumor to acquire the metastatic ability. During these five years the early diagnosis and prevention of deaths from pancreatic cancer could be fulfilled. Studies of Lennon *et al.* (2014), show that from the first mutation in the pancreas to the development of mutation sites, there could be a window of opportunity of detection of up to two decades. This is why, one of the most promising ways to improve the prognosis of pancreatic cancer is to develop effective early-detection strategies (Zhou *et al.*, 2017).

Nevertheless, current methods are not sufficient to improve the early-diagnosis (Rulyak *et al.*, 2003). The current diagnostic methods include, Computed Tomography (CT), Magnetic Resonance Imaging (MRI) and invasive endoscopic techniques

(Bunger *et al.*, 2011), which are not efficient nor comfortable for the patient. Especially in comparison over plasmatic biomarkers, for which you only need a blood sample (which remain the most promising approaches to improving long-term survival (Llop *et al.*, 2018).

## 1.2 Epidemiology

According to the World Health Organization (2019) “Epidemiology is the study of the distribution and determinants of health-related states or events (including disease), and the application of this study to the control of diseases and other health problems.” It is therefore a basic science of public health that deals with incidence, distribution and possible control of diseases, among others.

Pancreatic cancer is a disease occurring almost three times more frequently in developed countries (figure 4), as is the case of Europe, Asia and Northern and South America (Ferlay *et al.*, 2010).




Figure 4. Estimated age-standardized incidence rates worldwide in 2018, for pancreas cancer

(adapted from Word Health Organization, IARC, 2018)

Incidence and mortality rates have been increasing in Europe (Bosetti *et al.*, 2013), and North America (Kohler *et al.*, 2015).

Seeing specifically where most of the new cases are estimated to have occurred in 2018 (figure 5), Asia stands out, making 46,7% of all new cases worldwide. Europe comes in second place (before North America) with 28.9% (Globocan, 2018).




Figure 5. Estimated number of new cases of pancreatic cancer in 2018

(adapted from Globocan, 2018, <http://gco.iarc.fr>)

In 2018, despite not being one of the cancers with more new cases, pancreatic cancer counts to one of the eight with more deaths related to it, namely 4,6% of all cancer-related deaths (figure 6).




Figure 6. Number of new cases and deaths of cancer worldwide

(adapted from Globocan, 2018, <http://gco.iarc.fr>)

Comparing the number of incident cases and deaths in Europe and North America for Pancreatic Cancer (figure 7), the estimated age-standardized incidence is similar in both continents in 2018 (7.7 in Europe to 7.6 in North America) (Globocan, 2018).



Most pancreatic cancer patients are diagnosed at stage III or IV (79%), with only 21% having an early-diagnosis (stage I and II) (National Cancer Registration and Analysis Service, 2016, ISD Scotland, 2016).




Figure 7. Estimated number of new cases and deaths in 2018, comparing Europe and America

(Globocan, 2018, <http://gco.iarc.fr>)

## 1.3 Etiology

Tumorigenesis is the progress of the cancer cells into a clinically significant disease, taking years. It is a complex process which can have innumerable causes and may involve lifestyle, namely diet and physical activity, environmental factors and host factors, like inheritance or epigenetic changes (figure 8). As for the host factors they can influence the probability of cancer development, with the accumulation of genetic damage over time. The interaction of these three types of factors over time is critical to the development (or not) of a malignancy (Annex 1).




Figure 8. Factor affecting cancer process

(adapted from World Cancer Research Fund, 2018)

### 1.3.1 Non-modifiable risk factors

Pancreatic cancer, like all cancers and many other diseases, has a higher risk of occurring more frequently with **age** (World Cancer Research Fund, 2018). The risk increases especially after 65 years, with most diagnoses being made between the ages of 60 and 80 (Ferlay *et al.*, 2012).

In addition to that, it also varies with **race**, as the black population has an increased risk by 1.5.

Regarding pancreatic cancer in specific, it has the highest incidence of **venous thromboembolism** in the first year of diagnosis, in comparison to the other neoplasms

(Nakchbandi *et al.*, 2008). Patients with pancreatic cancer and venous thromboembolism also have a higher risk of poor prognosis and recurrence (Kondo *et al.*, 2018) (table 2).

According to Klein *et al.* (2018), individuals with **non-0 blood group** also have an increased risk of pancreatic cancer.

**Positive family history and some inherited genetic disorders** also enhance the risk of pancreatic cancer, as are the Lynch syndrome, Peutz-Jeghers syndrome, familial atypical multiple mole melanoma syndrome, among others (Greer *et al.*, 2007). Mutations in specific genes also increase the risk of developing this type of cancer (gene PRSS1, K-ras, p16, p53, BRCA2, ATM) (Slebos *et al.*, 2000).

An established risk factor is also the history of **previous pancreatic diseases**, as the case of pancreatitis. Although a correlation of higher risk can be made, most patients with pancreatitis do not develop cancer (Zheng *et al.*, 2013).

Considering that the **adult obtained height**, has as cause the number of cell divisions during fetal life and childhood, health and nutrition in childhood and age of sexual maturity, which changes hormonal microenvironment producing alterations in the level of growth factors (higher circulating levels of IGF-1 (Gunnell *et al.*, 2001, Bray *et al.*, 2006). Tall people have undergone more cell divisions, so there can be a higher potential for error during DNA replication, which can result in cancer (Le Roith *et al.*, 2001) (table 2).

Table 2. Summary non-modifiable risk factors

Risk factor	Reference
<b>Age (&gt; 65 years)</b>	Ferlay <i>et al.</i> , 2012, World Cancer Research Fund/American Institute for Cancer Research, 2018
<b>Black population</b>	World Cancer Research Fund/American Institute for Cancer Research, 2018
<b>Venous thromboembolism</b>	Kondo <i>et al.</i> (2018)
<b>Non-0 blood group</b>	Klein <i>et al.</i> (2018)
<b>Positive family history, some inherited genetic disorders</b>	Slebos <i>et al.</i> , 2000, Greer <i>et al.</i> , 2007
<b>Previous pancreatic diseases</b>	Zheng <i>et al.</i> , 2013
<b>Bigger attained adult obtained height</b>	Le Roith <i>et al.</i> , 2001, World Cancer Research Fund/American Institute for Cancer Research, 2018

### 1.3.2 Modifiable risk factors

As already known, cigarette smoking (or chewing) is one of the biggest causes of cancer, especially lung cancer. **Tobacco** use is proven to be the leading cause of pancreatic cancer too (Secretan *et al.*, 2009). It is estimated that 20% of pancreatic cancers are also caused by tobacco (IARC, 2012). The risk is increased by 74% for current smokers and 20% for former ones. Former smokers have an equal risk as non-smokers after 20 years (Lowenfels *et al.*, 2006). According to Parkin *et al.* (2010), this risk may even be bigger. It is definitely the leading identified cause of pancreatic cancer. Tobacco and its smoke contain more than 7000 chemical compounds, many identified as carcinogens (World Cancer Report, 2014). These carcinogens contribute through many pathways to the development of the neoplasm: DNA binding, mutations, inflammation, oxidative stress and epigenetic changes.

Another cause of a major part of the pancreatic cancers is **body fatness** (especially abdominal) and **physical inactivity**. It also increases the risk of almost all gastrointestinal cancers. With an increase of body fatness, there can also occur **insulin resistance and diabetes** (Hursting *et al.*, 2003), influencing levels of many other circulating hormones. This is why new-onset diabetes can be an early sign of pancreatic cancer (World Cancer Report, 2014). Due to the fact that insulin resistance is increasing in obesity, the pancreas compensates with and increased insulin

production. This **hyperinsulinemia** (Calle *et al.*, 2004), is an environment propitious to carcinogenesis, discouraging apoptosis (World Cancer Research Fund, 2018).

Due to the fact that adipocytes (fat cells) produce pro-inflammatory factors (Calle *et al.*, 2004), most obese individuals have a chronic inflammation (Rexrode *et al.*, 2003), which in turn can also promote carcinogenesis (Loffreda *et al.*, 1998).

By contributing to weigh control and influencing the overall metabolic state (World Cancer Research Fund, 2018), regular physical activity has proven to reduce the risks, as the reduction of sweetened food/ beverages and the consumption of vegetables, fruits and whole grains. Physical activity has also proven immunomodulatory effects (improves innate and acquired immunity) and promoting in this way a diminished risk of carcinogenesis (McTiernan *et al.*, 2008, Fridenreich *et al.*, 2010). Exercise also decreases oxidative stress improving in this way DNA repair mechanisms (Fridenreich *et al.*, 2010).

**Alcohol**, if consumed in abuse (more than three drinks a day) (World Cancer Research Fund, 2018), has a dose-response association (Baan *et al.*, 2009). Especially, but not only, ethanol is genotoxic contributing to carcinogenesis (World Cancer Report, 2014), being classified as a group 1 carcinogen (Duell *et al.*, 2012) firstly in 1988, by the IARC. It is thought that its metabolites, like acetaldehyde might have a higher carcinogenic potential (World Cancer Research Fund, 2018).

The vast majority of the pancreatic juice samples studied by Maekawa *et al.* (2018) had **bacterial DNA**, namely *Enterococcus faecalis*, suggesting that it may be involved in the progression from pancreatitis to cancer. In 2018, about 16% of the total cancers were due to **infections with viruses, bacteria and microparasites**, as *Helicobacter pylori*, hepatitis B and C viruses (World Cancer Report, 2014). Through compromising the immune system, infection with HIV could also increase the risk of virus-related viruses.

Exposure to **all types of radiation**, like ionizing, ultraviolet and electromagnet one, can increase the risk of malignancies (World Cancer Report, 2014). This is both true for natural (like the sun) and man-made sources.

General **air pollution** from vehicles, households and a wide range of industries and drinking-water contamination may have carcinogens, as is the case of aromatic hydrocarbons (HAP), benzenes, asbestos, nitrates and nitrites and arsenic (World Cancer Report, 2014).

Plastic is produced using **Bisphenol A**. This is the reason why more than 90% of USA, Europe and Asia are exposed and probably contaminated by it through food or liquids (Vandenberg *et al.*, 2010). In the 1980s it was established that Bisphenol A was not a robust carcinogen and recommended maximum daily safe dose was established to be

50 µg/kg/day ([www.epa.gov/iris/subst/0356.htm](http://www.epa.gov/iris/subst/0356.htm)). However, nowadays, with more than a hundred studies published since then, it is proven that this substance has an influence in the endocrine system (Keri *et al.*, 2007), causing changes in the cancer cells at concentrations within the limits of exposure (Vandenberg *et al.*, 2009). This evidence is however contested (Hengstler *et al.*, 2011).

**Pharmacological drugs**, including the ones used as antineoplastic agents in cancer therapy could also potentially induce cancer development (as prevent it) (World Cancer Report, 2014). Due to its unspecific genotoxicity, antineoplastic drugs could be responsible for inducing second cancers in supposed cured patients (table 3).

Concluding, many cancers can be prevented, by not engaging in those modifiable risk factors. The recommendations by the World Cancer Research Fund can be seen figure 9.




Figure 9. Cancer Prevention Recommendations

(adapted from World Cancer Research Fund, 2018)

Table 3. Modifiable risk factors

<b>Risk factor</b>	<b>Reference</b>
<b>Tobacco use</b>	Secretan <i>et al.</i> , 2009
<b>Body fatness, physical inactivity</b>	Hursting <i>et al.</i> , 2003
<b>Insulin resistance and diabetes</b>	Hursting <i>et al.</i> , 2003
<b>Alcohol</b>	World Cancer Research Fund/American Institute for Cancer Research, 2018
<b>Infections with viruses, bacteria and microparasites</b>	Maekawa <i>et al.</i> (2018), World Cancer Report, 2014
<b>All types of radiation</b>	World Cancer Report, 2014
<b>Air pollution</b>	World Cancer Report, 2014
<b>Bisphenol A</b>	Vandenberg <i>et al.</i> , 2009
<b>Pharmacological drugs</b>	World Cancer Report, 2014

## 1.4 Current Diagnosis Methods

Currently, the only way to diagnose asymptomatic pancreatic cancer is using imaging techniques (Winter *et al.*, 2006, Poruk *et al.*, 2013). Imaging the pancreas is also used for cancer staging, monitoring the response to treatment and detection of metastatic lesions.

Among others, the techniques used are computed tomography (CT), magnetic resonance imaging (MRI), positron emission tomography (PET), and endoscopic ultrasonography (EUS) (Morana *et al.*, 2010, Appel *et al.*, 2012, Fusaroli *et al.*, 2012, Raman *et al.*, 2012, Conrad *et al.*, 2013) (figure 10).




Figure 10. Algorithm for evaluation of a patient with suspicion of pancreatic cancer (adapted from Zhang *et al.*, 2018a)

#### 1.4.1 Multidetector computed tomography (MDCT)

For patients with suspicion of pancreatic adenocarcinoma MDCT is the most available and best-validated diagnostic method (Zhang *et al.*, 2018a). In addition to that it also the cheapest method, safe and non-invasive (Ahn *et al.*, 2009). It takes an image in which the disease and the rest of the pancreatic tissue can be well distinguished, as well as examination of distant disease (Brennan *et al.*, 2007). For these reasons, it is normally the first method of diagnosis being used (National Comprehensive Cancer Network, 2017). Nevertheless, it is a nephrotoxic method which exposes the patient to radiation (Al-Hawary *et al.*, 2014).

#### 1.4.2 Magnetic Resonance Imaging (MRI)

MRI has a superior imaging, allowing theoretically a diagnosis in earlier stage, which has proven to be not true in the practical case (with the same sensitivity as CT scanning) (Treadwell *et al.*, 2016). It is radiation free, but expensive and not possible if metal implants are present. For these reasons it normally is not the first choice (National Comprehensive Cancer Network, 2017).



### **1.4.3 Endoscopic Ultrasonography with fine needle aspiration (EUS)**

EUS consists on an upper gastrointestinal endoscopic examination with an echoendoscope positioned in the stomach near the pancreas under sedation (Zhang *et al.*, 2018a). It is a safe, less invasive diagnosis method, having high sensitivity and detecting small lesions (Puli *et al.*, 2013). It also allows the collection of a sample for biopsy (by needle aspiration) (Hewitt *et al.*, 2012).

However, it is not always available, is operator dependent and it does not allow the detection of distant metastasis (Chen *et al.*, 2012).

### **1.4.4 Positron Emission tomography (PET)**

PET consists on enhancing the metabolism of glucose in cancer cells. These overexpress glucose transporter 1 and can accumulate glucose like this (Llop *et al.*, 2018). For detection, Fluorine 18-fluorodeoxyglucose (FDG) is used. FDG is a glucose analogue, used as a radiotracer. FDG-PET has yet a low sensitivity and specificity, resulting often in false-negatives (due to hyperglycemia) and false-positives (as FDG levels increase in inflammations for example) (Matsumoto *et al.*, 2013). PET has been showing to have better, more sensitive results for monitoring response to treatment in comparison to CT and for detecting recurrence (Sperti *et al.*, 2010, Kinupe *et al.*, 2017). Due to its wide anatomic coverage it is easier for metastatic detection in the entire body (Dibble *et al.* 2012, Lee *et al.*, 2014).

### **1.4.5 Biopsy**

Confirmation procedure, consisting on removing a small sample of tissue to be examined under microscope. This can occur by fine-needle aspiration (inserting a needle into the pancreas) or during EUS.

### **1.4.6 Blood test**

The best, most invasive, easier and quicker way to diagnose is by a simple blood test, looking for biomarkers. Biomarkers can be used with different goals, namely early detection, therapeutic monitoring, follow-up after surgery and to guide treatment decisions (Llop *et al.*, 2018).

For pancreatic cancer the only used is CA19-9, with no diagnostic goal though. It is currently FDA-approved for monitoring the disease.

## **1.5 Therapeutics**

Currently the only curative option is surgery, which can only be performed in early-stage pancreatic cancer (Costello *et al.*, 2012).

Unlike in most cancers, cytotoxic agents are still first-line treatment for pancreatic cancer (Shi *et al.*, 2018). Nevertheless, there are not many pharmacological agents that can be used and therefore do not vary much with the type of pancreatic cancer.

According to Shi *et al.* (2018), the levels of CA19-9 should be monitored while choosing the therapeutic for patients with advanced pancreatic cancer.

### **1.5.1 Surgery**

The only option that actually heals the patient. It can consist in removing only the part of the pancreas where the tumor is located or the entire pancreas. A person without pancreas can live normally needing insulin and enzyme replacement.

### **1.5.2 Chemotherapy**

Consists in the use of toxic pharmacological substances injected or taken orally isolated or in combination with radiation therapy (Shi *et al.*, 2018). In combination it is normally used in cases of wide spread pancreatic cancers or in cases where the tumor has first to be shrunken so that surgery is possible. It is also possible after surgery to prevent recurrence of the tumor.

### **1.5.3 Radiation Therapy**

Cancer cells are destroyed with the use of highly energetic radiation, such as X-rays and protons. As chemotherapy, it can be used when surgery is not an option, to reduce the tumor before surgery or after surgery. Often in combination with chemotherapy.

## **2 Pancreatic Carcinogenesis**

### **2.1 Pathology**

Tumor mass consists of pancreatic stellate cells, immune cells, lymphatic and vascular endothelial cells, pathologically increased nerves and extracellular matrix (Duan *et al.*, 2017).

The infiltrating ductal adenocarcinoma deduces an intense desmoplastic stromal reaction (Li *et al.*, 2012, Rahib *et al.*, 2014), creating a very complex tumor

microenvironment (Jiang *et al.*, 2018). This complexity explains the lethality of this cancer, promoting its development (O'Neil *et al.*, 2012, Duan *et al.*, 2017). Due to its complexity, biopsies may mislead to false negative results (Olive *et al.*, 2009).

Desmoplasia is the growth of fibrous or connective tissue. Although it may occur in benign circumstances like in scar tissue after surgery, it normally is associated with malignant neoplasms (causing dense fibrosis around it and invading healthy tissues like this) (Jiang *et al.*, 2018).

Pancreatic stellate cells are thought to be the most responsible for the desmoplasia (Masamune *et al.*, 2015). They occur normally in the pancreas, but they can be activated when in contact with pancreatic cells, synthesizing biologically active molecules (Pothula *et al.*, 2016). They can also inhibit apoptosis and promote stem cells phenotypes of pancreatic cancer cells (Erkan *et al.*, 2008), being therefore responsible for the resistance of pancreatic cancer to chemotherapy, distant metastasis and a poor prognosis (Hwang *et al.*, 2008).

## 2.2 Genetics

Around 10% pancreatic cancer is of familial origin, meaning that some inherited mutated genes increase the risk of its development (Shi *et al.*, 2009).

Exosome sequencing of ductal adenocarcinoma showed 16 significantly mutated genes (World Cancer Report, 2014). Including one oncogene (*KRAS*), three tumor suppressor genes (*CDKN2A*, *TP53*, *SMAD4*), *MLL3*, *ATM*, *TGFBR2*, *ARID1A*, and *SF3B1* (Jones *et al.*, 2008).

Biankin *et al.* (2012) also discovered novel mutated genes: genes involved in chromatin modification (*EPC1* and *ARID2*) and DNA damage repair, and other mechanisms (*ZIM2*, *MAP2K4*, *NALCN*, *SLC16A4*, and *MAGEA6*).

Studying the mutated genes, does not only help in monitoring high-risk populations, but also in the development of personalized therapy, like the poly (ADP-ribose) polymerase inhibitors or mitomycin C for cancers with *BRCA2* or *PALB2* mutations (World Cancer Report, 2014).

## 3 Current State of the Art in Diagnostics

### 3.1 Novelties in Pancreatic Biomarkers

There is no current reliable early-state diagnostic biomarker for pancreatic cancer (Zhang *et al.*, 2018b).

Due to the fact that most patients are only diagnosed with advanced pancreatic cancer, the samples to be studied for biomarkers come from them, making the identification of early-stage biomarkers more difficult (Resovi *et al.*, 2018). This is why the use of in vivo models is fundamental.

Biomarkers are the most promising early-diagnosis method, due to its low cost, convenience, quickness and minimal invasiveness (Llop *et al.*, 2018). However, it has been showed recently that no single biomarker could be reliable enough for cancer diagnosis, making a combination of biomarkers and incorporation of other clinical factors (as imaging techniques) a better approach (Capello *et al.*, 2017, Chang *et al.*, 2017). Figure 11 illustrates the potential early-stage pancreatic cancer biomarkers currently being studied.




Figure 11. Overview of circulating biomarkers for early detection of pancreatic cancer

(adapted from Zhang *et al.*, 2018b)

### 3.1.1 Traditional tumor biomarkers

#### 3.1.1.1 Carbohydrate Antigen 19-9 (CA19-9)

CA19-9 is a modified Lewis blood group carbohydrate antigen (Young *et al.*, 2018), called siayil Lewis A (SLe<sup>a</sup>). It is embedded on cell surface molecules, namely glycoproteins, gangliosides and mucins (Goh *et al.*, 2017, Young *et al.*, 2018), and is expressed in pancreatic and hepatobiliary diseases (Zhang *et al.*, 2018a). It is the only FDA approved biomarker for pancreatic cancer (Locker *et al.*, 2006), widely used with ELISA (Yue *et al.*, 2011) but with many limitations (Wong *et al.*, 2008).

It is used for monitoring the response of pancreatic cancer to treatment (Wong *et al.*, 2008, Yue *et al.*, 2011), as a prognostic biomarker (national cancer observatory) and for monitoring recurrence after surgical resection (Ballehaninna *et al.*, 2012) having a lack of sensitivity and specificity for early-diagnosis (Huang *et al.*, 2014). Their values to diagnosing pancreatic cancer vary according to different authors: 82 and 90% respectively (Llop *et al.*, 2018) and according to Huang *et al.* (2014), it has a variable sensitivity of ~85% and specificity of ~85%. It has been shown that CA19-9 is upregulated up to two years before the diagnosis of pancreatic cancer, showing its good potential (O'Brien *et al.*, 2015).

According to a study of Haab *et al.* (2015), most studies performed with CA19-9 do not use the same standards, varying from institution to institution. Feng *et al.* (2013), performed a large cohort, using the same rigorous conditions in every institution and key control groups. The goal was to understand if the different results obtained until that point, had divergent results due to the use of different standards. For this study they compared two antibody-based assay kits. As a global result, they reached similar AUC of 0.77 (area under the curve, it measures the accuracy with an AUC of 1 representing a perfect test) and comparable to the ones obtained in previous studies. However, these results varied a lot between patients, showing here a potential to be farther studied (Young *et al.*, 2018).

As a prognostic biomarker it performs better in symptomatic patients (Ballehaninna *et al.*, 2012, Huang *et al.*, 2014). Nevertheless, it is not reliable when used alone (Singh *et al.*, 2011).

According to Llop *et al.* (2018), it is not expressed in 10-20% of the Lewis antigen-negative population, due to a genetic deficiency of fucosyltransferase activity (Capello

*et al.*, 2017). In general, about 5-10% of the whole population does not express it (Herrerros-Villanueva *et al.*, 2016, Root *et al.*, 2018).

In addition to that, it can lead to false-negatives, as only 65% of patients with pancreatic cancer have a high expression of CA19-9 (Gold *et al.*, 2013).

It can also be falsely positive when other pathologies are present, like biliary infection, inflammation or obstruction, chronic pancreatitis and obstructive jaundice (Marrelli *et al.*, 2009), hepatic and pancreatic cysts (Llop *et al.*, 2018), other benign gastrointestinal conditions (Locker *et al.*, 2006) or other cancers, like colorectal and breast (Llop *et al.*, 2018). It has also racial and sex expression variations, being highest in Caucasians (Resovi *et al.*, 2018).

In order to overcome these limitations and improve efficiency a combination of CA19-9 and other biomarkers has good potential (Faca *et al.*, 2008, Chang *et al.*, 2009, Brand *et al.*, 2011 Gold *et al.*, 2013, Lennon *et al.*, 2014).

Many of these combinations with CA19-9 are being studied. The combination with **albumin and IGF** (Insulin-like growth factor 1) (Goh *et al.*, 2017), can distinguish between pancreatic cancer and chronic pancreatitis with a high sensibility (93,6%) and high specificity (95%) (Park *et al.*, 2012, Ferri *et al.*, 2016).

Makawita *et al.* (2013), studied regenerating islet-derived 1 beta (REG1B), syncollin (SYCN), anterior gradient homolog 2 protein (AGR2), and lysyl oxidase-like 2 (LOXL2). The combination between **CA19-9, SYNC and REG1B** had the best results, with an AUC of 0.9.

In combination with **ICAM-1** (Intercellular Adhesion Molecule 1) and **OPG** (Osteoprotegerin), Brand *et al.* (2011), reported a sensitivity for diagnosis from 88% and a specificity of 94%. It is a highly selective combination that had negative results from almost all breast, lung and colorectal cancers.

In combination with **cathepsin D and MMP-7** (matrix metalloproteinase-7) it even has a higher AUC (Park *et al.*, 2012).

Zeh *et al.* (2005) reported a combination of CA19-9 and serum cytokines (**HGF, MCP-1, IP-10, Eotaxin**) with very good results: sensitivity 85.7% and specificity of 92.3% to distinguish PC from healthy controls and sensitivity of 98%, specificity of 96.4% to distinguish PC from chronic pancreatitis. CA19-9 in combination with **albumin, C-reactive protein and interleukin 28** demonstrated sensitivity was 99.39% for all-stages, 96.10% for early-stage and 98.80% for advanced-stage pancreatic cancer at 90% specificity when discriminating between cancer and healthy individuals.

Cohen *et al.* (2017), studied the combination of four plasma proteins (**CA19-9, CEA** (Carcinoembryonic antigen), **HGF** (Hepatocyte growth factor), **OPN** (osteopontin)) with ctDNA testing for **KRAS mutations**. Although it outperformed

CA19-9 alone, it had a sensitivity of only 64%, making an optimization of the assays crucial to see if it could be an option to test furtherly.

Capello *et al.* (2017) researched already studied protein candidates to come with the best combination, namely **TIMP1 (TIMP metalloproteinase inhibitor 1)**, **LRG1 (Leucine-rich alpha-2-glycoprotein 1)** and **CA19-9**, which outperformed CA19-9.

Combining CA19-9 with MUC5AC (Mucin 5AC), which in literature seems to have very good biomarker properties (is over-expressed in pre-cancerous lesions), Kaur *et al.* (2017), found it to have a better performance in ELISA.

The same was proven by Kim *et al.* (2017) for the combination **thrombospondin-2 (THBS2)** and **CA19-9**.

Sefrioui *et al.* (2017) studied a panel consisting in **CA19-9, ctDNA and CTC analysis**. In comparison with EUS-FNA alone it increased significantly the sensitivity and specificity.

According to Llop *et al.* (2018), combining **CA19-9 with miR-16 and miR196-a** it improves its performance to an AUC from 0,98.

All these combinations have still to be correctly validated first (table 4).

Table 4. Summary of CA19-9 combinations under study as early-stage diagnostic pancreatic cancer biomarkers

Biomarker Panel (concentration)	AUC, sensitivity, specificity	Reference
<b>Carbohydrate Antigen 19-9 (CA 19-9)</b>	AUC: 0.77; Sensitivity: ~85%; Specificity: ~85%	Feng <i>et al.</i> (2013) Huang <i>et al.</i> , 2014
<b>CA 19-9, albumin and IGF</b>	Sensitivity: 93,6%; Specificity: 95%	Goh <i>et al.</i> , 2017 Park <i>et al.</i> , 2012 Ferri <i>et al.</i> , 2016
<b>CA 19-9, SYNC, REG1B</b>	AUC: 0.9;	Makawita <i>et al.</i> , 2013
<b>CA 19-9, ICAM-1, OPG</b>	Sensitivity: 78%; Specificity: 94%	Brand <i>et al.</i> , 2011
<b>CA 19-9, cathepsin D, MMP-7</b>	AUC: 0.9 Sensitivity: 88%; Specificity: 80%	Park <i>et al.</i> , 2012
<b>CA 19-9, HGF, MCP-1, IP-10, Eotaxin</b>	Sensitivity: 85.7%; Specificity: 92.3%	Zeh <i>et al.</i> , 2005
<b>CA 19-9, albumin, C-reactive protein and interleukin</b>	Sensitivity: 99.39%; Specificity: 90%	Zeh <i>et al.</i> , 2005
<b>CA19-9, CEA, HGF, OPN with ctDNA testing for KRAS mutations</b>	Sensitivity: 64%; Specificity: 99.5%	Cohen <i>et al.</i> , 2017
<b>CA19-9, TIMP1, LRG1</b>	AUC: 0.95; Sensitivity: 75%; Specificity: 95%	Capello <i>et al.</i> , 2017
<b>CA19-9, MUC5AC</b>	AUC: 0.87; Sensitivity: 83%; Specificity: 83%	Kaur <i>et al.</i> , 2017
<b>CA19-9, THBS2</b>	Sensitivity: 87%; Specificity: 98%	Kim <i>et al.</i> , 2017
<b>CA19-9, ctDNA and CTC analysis</b>	Sensitivity: 78%; Specificity: 91%	Sefrioui <i>et al.</i> , 2017
<b>CA19-9, miR-16 and miR196-a</b>	AUC: 0.98	Llop <i>et al.</i> , 2018



### 3.1.1.2 Other conventional biomarkers

There are many other conventional biomarkers (Zhang *et al.*, 2018b), namely CA242, CA72-4, CA125, CEA (Bunger *et al.*, 2011). Although they have limited early-diagnostic potential (Zhang *et al.*, 2015), they are promising for Lewis-negative population (in which CA19-9 has no utility).

An example of this high specificity seen for Lewis-negative patients are CEA and CA125 (98% and 93,8% respectively) (Luo *et al.*, 2017).

However, this would imply first testing a possible patient regarding the state of the Lewis-group, which is not ideal

### 3.1.2 Metabolites

According to Hanahan *et al.* (2011), due to the fact that cancer cells can rewire metabolically, they can survive and proliferate even with oxygen and nutrient deficiency (called Warburg effect (Warburg *et al.*, 1956)). In the specific case of pancreatic cancer, these cells survive in hypoxia, desmoplasia (Jiang *et al.*, 2018) and hypovascularization.

A malignancy could therefore be diagnosed in the presence of aberrant low-molecular weight substances (Denkert *et al.*, 2012) deriving from this abnormal biochemical state (Vermeersch *et al.*, 2013).

Because of the different demands of pancreatic cancer cells, especially metabolic ones, they can rewire the metabolism. This happens normally with a mutation of the oncogene KRAS (which is normally present in pancreatic cancer (Biankin *et al.*, 2012)). These different metabolic demands are supported by the fibroblasts, which mediate metabolite exchange (Ozdemir *et al.*, 2014, Rhim *et al.*, 2014) and pancreatic stellate cells (Sousa *et al.*, 2016).

Alterations in the metabolism include alterations in metabolic enzyme, accumulation of intermediates (Zhou *et al.*, 2012), making amino acids, lipids, fatty acids and bile acids in serum potential biomarkers (Kobayashi *et al.*, 2013, Ritchie *et al.*, 2013, Zhang *et al.*, 2013). One amino acid who is being studied is palmitic acid in serum. According to Di Gangi *et al.* (2016), it has been showed to have better results in early-diagnosis of pancreatic cancer than CA19-9 with an AUC of 1.0.

Mayers *et al.* (2014), also studied plasmatic amino acids, coming to the conclusion that an increase deriving from muscle catabolism could increase the risk of developing pancreatic cancer (Young *et al.*, 2018).

Four other serum metabolites showed a higher sensitivity and lower false-negative rate when comparing with CEA and CA19-9, namely xylitol, 1,5-anhydro-D-glucitol, histidine, and inositol (Sakai *et al.*, 2016).

There are extreme changes during different stages of pancreatic cancer. Metabolomics is therefore a sensitive indicator for monitoring pre-cancerous lesions and diagnosing early-stage (Yuan *et al.*, 2016). For example, Kynurenate and methionine levels are only upregulated in pre-cancerous lesions and decreased in cancer.

Yuan *et al.* (2016) studied 82 metabolites in a prospective cohort study, concluding that two metabolites (isocitrate and aconitate) were associated with survival of pancreatic cancer patients. Both of these metabolites are intermediates of the tricarboxylic acid cycle. In addition to that, inherited ACO1 genotypes also influenced survival of patients.

Nowadays there have been identified already 50 differences in serum metabolites between pre-cancerous lesions and pancreatic cancer, making metabolites a good potential panel for biomarkers (LaConti *et al.*, 2015).

Different enzymes and intermediates influence the concentration of a single metabolite, making the development of a panel with a combination of biomarkers crucial for a good diagnostic.

As for other potential biomarkers, they still have to be studied in larger populations and under standardized conditions (Zhang *et al.*, 2018b) (table 5).

Table 5. Principal Metabolites being studied as early-stage diagnostic pancreatic cancer biomarkers

Biomarker Panel	AUC, sensitivity, specificity	Reference
<b>Palmitic acid</b>	AUC: 1.0; Sensitivity: 100%; Specificity: 100%	Di Gangi <i>et al.</i> , 2016
<b>Xylitol, 1,5-anhydro-D-glucitol, histidine, and inositol</b>	Sensitivity: 84.1%; Specificity: 84.1%	Kobayashi <i>et al.</i> , 2013

### 3.1.3 Cell-free DNA

By apoptosis or necrosis nucleic acids can be freed into the extracellular environment, originating cell-free DNA (cfDNA) (Zhang *et al.*, 2018b). Cui *et al.* (2008), showed that patients with solid tumors normally have elevated levels of cfDNA.

Circulating tumor DNA (ctDNA) carries mutations corresponding to the somatic mutations in the primary tumor (Kinugasa *et al.*, 2015), and it is believed that it correlates to its burden (Fleischhacker *et al.*, 2007, Diaz *et al.*, 2014, Newman *et al.*, 2014), meaning that analyzing ctDNA from liquid biopsies could indicate tumor state and genetics. Bettegowda *et al.* (2014), showed that in stage I to III of pancreatic cancer, about 40% of patients have detectable ctDNA and about 90% for stage IV tumors.

Liquid biopsies are one of the most recent approaches in oncology, but still not suitable to replace tissue biopsies (Imamura *et al.*, 2016, Pishvaian *et al.*, 2016, Riva *et al.*, 2016).

### **3.1.3.1 Somatic mutations**

In pancreatic cancer, inactivating mutations in tumor suppressor genes, like CDKN2A, TP53, SMAD4, BRCA2 are found mostly in late states. In contrast to this, the most predominant genetic characteristics are KRAS mutations (Zhang *et al.*, 2018b). They occur even in premalignant lesions and in high rate, being detected in serum, pancreatic juice and feces (Jones *et al.*, 2008). However, it is not specific for pancreatic cancer as it also occurs in other diseases like chronic pancreatitis (Yanagisawa *et al.*, 1993).

Its importance consists in controlling patients with pancreatic intraepithelial neoplasia. Almost 90% of these patients have a KRAS mutation with its rate correlating to the grade of the disease implying that a KRAS mutation is an early event during tumorigenesis (Kanda *et al.*, 2012). There was also found a relation between KRAS mutations and clinical stage (Castells *et al.*, 1999).

This potential biomarker still needs to be furtherly studied, as results of current studies do not always match (Mullcahy *et al.*, 1998, Maire *et al.*, 2002).

The detection rates vary from 35% of plasma samples of pancreatic cancer patients (Uemura *et al.*, 2004) and 48% (Bettegowda *et al.*, 2014) to 63% of pancreatic patients (Kinugasa *et al.*, 2015).

Mutations in TP53 and SMAD4, in comparison with KRAS mutations, do not occur very often (Zhang *et al.*, 2012). This is the reason why there are only a few studies on them. These mutations occur normally later, making them not suitable for early-stage diagnosis (Maitra *et al.*, 2003) (table 6).

Table 6. Somatic Mutations of Cell-free DNA as early-stage diagnostic biomarkers

Biomarker Panel	AUC, sensitivity, specificity	Reference
<b>KRAS mutation</b>	Detection rate: 35-63% of pancreatic cancer patients	Uemura et al., 2004 Kinugasa et al., 2015
<b>TP53 and SMAD4 mutations</b>	Lower detection rates than KRAS mutation	Zhang et al., 2012

### 3.1.3.2 Epigenetic modulations

Epigenetic modifications of cfDNA (cell-free DNA), especially alterations in methylation pattern are very common in cancer, as it is the case of hypermethylation of tumor genomic DNA and hypermethylation of tumor suppressor genes (Zhang *et al.*, 2018b). Sato *et al.* (2003), discovered several targets of abnormal DNA methylation in pancreatic cancer. UCHL1, NPTX2, SARP2, CLDN5, FOXE1, CDH3.

CD1D, KCNK12, CLEC11A, NDRG4, IKZF1, PKRCB and KRAS are currently under clinical study after resulting in 75% sensitivity and 95% specificity (Kisiel *et al.*, 2015). Other methylation biomarkers associated to the disease are ppENK, cyclin D2, sparc-7, Osteonectin and TSLC1. There has not been found further validation and clinical application for them yet (Fukushima *et al.*, 2003, Matsubayashi *et al.*, 2003, Sato *et al.*, 2003).

There are, in addition to that, abnormal methylation profiles in specific cfDNA regions. Although P16 and proproenkephalin promoters are hypermethylated in plasmatic DNA, their detection rates are from about 30% and 25% accordingly (Jiao *et al.*, 2007).

After analyzing 30 plasma samples, Liggett *et al.* (2010), combined some targets, particularly 14 gene promoters differentiating chronic pancreatitis patients from controls (sensitivity of 81,7% and specificity of 78%) and from pancreatic cancer patients (sensitivity of 91,2% and specificity of 90,8%) according to its methylation status. The panel included, CCND2 (cyclin D2), DAPK1 (death-associated protein kinase 1), ESR1 promA (estrogen receptor 1 promoter A), HMLH1 (human mutL homolog 1), MGMT (O-6-methylguanine-DNA methyltransferase), MUC2, (mucin 2, oligomeric mucus/gel-forming), MYOD1 (myogenic differentiation 1), CDKN2B (cyclin-dependent kinase inhibitor 2B), CDKN1C (cyclin-dependent kinase inhibitor 1C), PGK1 (phosphoglycerate kinase 1), PGR-proximal (pro-gesterone receptor proximal promoter), RARb (retinoic acid receptor beta), RB1 (retinoblastoma 1), SYK (spleen tyrosine kinase).

A combination of CCND2, SOCS1 and THBS1 has also a very good potential for early-stage diagnosis, with a sensitivity of 76% and a specificity of 59% (Melnikov *et al.*, 2009).

Studying the methylation status of BNC1 (Basonuclin 1) and ADAMTS1 (ADAM Metalloproteinase with Thrombospondin Type 1 Motif 1) in cfDNA could also be used for early-stage detection (81% sensitivity, 85% specificity) (Yi *et al.*, 2013).

Still, epigenetic modifications need to be furtherly studied but have a very good potential. The incidence of aberrant DNA methylation at select cpg islands higher than incidence of genetic mutations and they have fewer false-negatives. This aberrant epigenetic alteration is an early event during tumorigenesis, it leads to gain/loss of function of critical molecules in cancer cells and the DNA methylation status is stable, making it easily detected with great sensitivity, even with contamination (Zhang *et al.*, 2018b) (table 7).

Table 7. Principal methylation patterns of gene promoters in cfDNA as biomarkers

Biomarker Panel (Methylation status of gene promoters in cfDNA measured)	AUC, sensitivity, specificity	Reference
<b>UCHL1, NPTX2, SARP2, CLDN5, FOXE1, CDH3, CD1D, KCNK12, CLEC11A, NDRG4, IKZF1, PKRCB and KRAS</b>	Sensitivity: 75% Specificity: 95%	Kisiel et al., 2015
<b>ppENK, cyclin D2, sparc-7, Osteonectin and TSLC1</b>	With no validation yet	Fukushima et al., 2003, Ueki et al., 2002, Matsubayashi et al., 2003, Sato et al., 2003
<b>CCND2, DAPK1, ESR1 promA, HMLH1, MGMT, MUC2, MYOD1, CDKN2B, CDKN1C, PGK1, PGR-proximal, RARb, RB1, SYK</b>	Sensitivity: 91,2% Specificity: 90,8%	Liggett et al., 2010
<b>CCND2, SOCS1, THBS1</b>	Sensitivity: 76% Specificity: 59%	Melnikov et al., 2009
<b>BNC1, ADAMTS1</b>	Sensitivity: 81%; Specificity: 85%	Yi et al., 2013

### 3.1.4 Autoantibodies

Cancer patients normally develop an immune system dysfunction (Zhang *et al.*, 2018b). Though the immune response is not strong enough to have clinical manifestations (Kobold *et al.*, 2010), circulating autoantibodies could be potential biomarkers for diagnosing cancer.

In some types of cancer, including pancreatic, there are produced autoantibodies against tumor-associated antigens, ie. misfolded, overexpressed, aberrantly modified, ectopically expressed and mutated tumor proteins, (Desmetz *et al.*, 2011, Kaur *et al.*, 2012). These autoantibodies can indirectly reflect altered genetics and proteomics but can only be detected in low frequency (Zhang *et al.*, 2018b). Due to tumor heterogeneity, detection is even more difficult.

Despite the fact that, in a study by Bracci *et al.* (2012), there were analyzed several autoantibodies, concluding that there are significant differences in expression levels in pancreatic cancer patients, their diagnostic value was poor with an AUC < 0.7.

Recently, the main focus on autoantibodies research has been Anti-mucin 1 antibodies (MUC1). MUC1 is a membrane associated glycoprotein that is overexpressed in some cancers, including pancreatic one and is associated with CA19-9 (Zhang *et al.*, 2018b). In the last years, Gold *et al.* (2006), researched a monoclonal antibody against MUC1 with a sensitivity of 77% and a specificity of 95% for diagnosing pancreatic cancer.

Other autoantibodies being researched include autoantibodies to two acidic isoforms of glycolytic enzyme enolase (ENOA 1 e 2), which seem to be more frequent in patients with normal CA19-9 levels (Tomaino *et al.*, 2011), Ezrin (Capello *et al.*, 2013), vimentin isoform (Hong *et al.*, 2006), and calreticulum isoforms (Hong *et al.*, 2004) (table 8).

Neoantigen are new immunogenic protein sequences produced by malignancies. They are absent from the normal human genome and are primarily created by tumor-specific mutations in the genome. They could be potential targets for diagnoses and immunotherapy (Schumacher *et al.*, 2015).

Table 8. Autoantibodies with potential as biomarkers

Biomarker Panel	AUC, sensitivity, specificity	Reference
<b>Monoclonal antibody against MUC1</b>	Sensitivity: 77% Specificity: 95%	Gold et al., 2006
<b>ENOA 1 e 2</b>	Sensitivity: 62% Specificity: 97%	Tomaino et al., 2011
<b>Ezrin</b>	AUC: 0.9 Sensitivity: 93.2% Specificity: 75.5%	Capello et al., 2013
<b>Vimentin isoform</b>	With no validation yet	Hong et al., 2006
<b>Calreticulum isoforms</b>	With no validation yet	Hong et al., 2004

### 3.1.5 Cell-free noncoding RNA (ncRNA)

Genes code proteins, but only about 2% of all genes. At least 75% of them do not encode anything, being called noncoding RNA (ncRNA) (Djebali *et al.*, 2012). ncRNA include microRNA, small interfering RNA, piwi-interacting RNA, small Cajal body-specific RNA and long noncoding RNA (St Laurent *et al.*, 2015).

It is suspected that ncRNA play regulatory roles, modifying gene expression at multiple levels through interactions with DNA, RNA and proteins during physiological processes and tumor development (Presner *et al.*, 2011), especially miRNA and lncRNA that are specific of pancreatic cancer (Muller *et al.*, 2015).

They can be detected in circulation, being potential biomarkers (Zhang *et al.*, 2018b).

### **3.1.5.1 microRNA (miRNA)**

miRNAs are small non-coding RNAs (18-25 nucleotides) that bind to complementary mRNA and inhibit gene expression. They regulate expression of more than 30% of human genes and may also function as tumor promoters or suppressors (Zhang *et al.*, 2018b) as they can regulate gene expression. In oncogenesis there occurs a deregulation for which miRNAs may be responsible (Iorio *et al.*, 2009, Li *et al.*, 2013). Alterations in miRNA expression can, therefore, drive to malignancies (Lu *et al.*, 2005). On the blood, miRNA can be found circulating as free RNA attached to hAgo2 (an argonaute protein, that pairs to miRNA silencing it) (Fujita *et al.*, 2004, Wakatsuki *et al.*, 2005), and incorporated in exosomes (that protect it from degradation (Valadi *et al.*, 2007)).

Due to their small size, stability, easy detection and convenient extraction they promise to be good biomarkers and to be furtherly studied. They are also deregulated in pancreatic diseases, making it able to discriminate pancreatic cancer from other malignancies (Bauer *et al.*, 2012, Schultz *et al.*, 2014).

Nonetheless, there is still no accepted internal control, studies lack a normalized standard and they can be bound to proteins or integrated in vesicles making their accurate quantification difficult. miRNA profiles are also very dynamic and easily and constantly influenced by other pathologies. Their origin and mechanisms are too still unclear (Zhang *et al.*, 2018b). Because of these, developing better techniques for its evaluation could be crucial.

It was recently shown that miR-10, miR-21, miR-22, miR-155 when expressed in different rates could indicate pancreatic cancer (Roldo *et al.*, 2006, Bloomston *et al.*, 2007, Lee *et al.*, 2007). They could have a higher diagnostic value than CA19-9.

According to a study performed by Schultz *et al.* (2014), where there were studied over 700 miRNAs, they found miR-145, miR-150, miR-223, miR-636 to be dysregulated in this malignancy. However, they proved to have an AUC (0.93), sensibility (0.85) and specificity (0.85) no superior than CA19-9 (AUC 0.9).

Another big, multicenter study focused on miR-486-5p (Xu *et al.*, 2016), coming also to results comparable with the diagnostic value of CA19-9. Cao *et al.* (2016) also used



the potential of miR-486-5p. These authors tested two different panels containing this miRNA, namely miR-486-5p, miR-126-3p, miR-106b-3p (AUC: 0.89; sensitivity: 82.7%; specificity: 84.4%) and miR-486-5p, miR-126-3p, miR-106b-3p, miR-938, miR-26b-3p, miR-1285 (AUC: 0.89; sensitivity: 82.3%; specificity: 81.4%).

Li *et al.* (2013), evaluated also over 700 circulating miRNAs. miR-1290 was the one with the best results, with an AUC of 0.96 for differentiating controls from pancreatic cancer patients (performing better than CA19-9) (Li *et al.*, 2013).

Continuing to study and identifying altered levels of miRNA on the precursor lesions of pancreatic cancer could provide further comprehension of these processes (du Rieu *et al.*, 2010, Xue *et al.*, 2013). Also, miR-21 (du Rieu *et al.*, 2010, Caponi *et al.*, 2013), miR-155 (Ryu *et al.*, 2010, Caponi *et al.*, 2013), miR-196 (Slater *et al.*, 2014), and miR-210 (Xue *et al.*, 2013), are overexpressed in these precancerous lesions in tissue, serum, cyst fluid and stool making them promising biomarkers to be studied (Hernandez *et al.*, 2016).

Schultz *et al.* (2014), identified two different miRNA panels that can differentiate healthy controls from pancreatic cancer patients. The first one consists of ten miRNAs (miR-26b, miR-34a, miR-122, miR-126, miR-145, miR-150, miR-223, miR-505, miR-636, miR-885.5p) and CA19-9. The second one includes four biomarkers, namely miR-145, miR-150, miR-223 and miR-636. For this second panel, the results were superior than those for CA19-9 alone (AUC: 0.93). Furthermore, it is also established that miR-216 and miR-217 are downregulated and miR-143, miR-145, miR-146, miR-148, miR-150, miR-155, miR-196a, miR-196b, miR-210, miR-222, miR-223 and miR-31 are up-regulated in pancreatic cancer (Bloomston *et al.*, 2007, Szafranska *et al.*, 2007, Hanoun *et al.*, 2010, Liffers *et al.*, 2011).

miR-223, needing still further studying, might be indicated for early-stage diagnostic and predicting malignant potential from pre-cancerous lesions (Komatsu *et al.*, 2015).

There are still very few studies associating miRNA levels and preneoplastic conditions (Komatsu *et al.*, 2015). Studying patients with PanIN (pancreatic intraepithelial neoplasia) and IPMN (intraductal papillary mucinous neoplasm of the pancreas) and following them could help understand it.

That is why miRNAs are currently one of the most promising and most studied biomarkers for early-pancreatic cancer detection (Herreros-Villanueva *et al.*, 2016). There is still a larger validation needed (table 9).

Table 9. Summary, miRNA as potential biomarkers

Biomarker Panel (concentration)	AUC, sensitivity, specificity	Reference
<b>CA19-9, miR-16 and miR196-a</b>	AUC: 0.98	Llop et al. (2018),
<b>CA19-9, miR-145, miR-150, miR-223, miR-636</b>	AUC: 0.93; Sensitivity: 85%; Specificity: 85%	Schultz et al., 2014
<b>miR-1290</b>	AUC: 0.96	Li et al., 2013
<b>CA19-9, miR-26b, miR-34a, miR-122, miR-126, miR-145, miR-150, miR-223, miR-505, miR-636, miR-885.5p</b>	AUC: 0.93; Sensitivity: 85%; Specificity: 85%	Schultz et al., 2014
<b>miR-486-5p, miR-126-3p, miR-106b-3p</b>	AUC: 0.89; Sensitivity: 82.7%; Specificity: 84.4%	Cao et al., 2016
<b>miR-486-5p, miR-126-3p, miR-106b-3p, miR-938, miR-26b-3p, miR-1285</b>	AUC: 0.89; Sensitivity: 82.3%; Specificity: 81.4%	Cao et al., 2016

### 3.1.5.2 Long non-coding RNA (lncRNA)

lncRNAs consist in more than 200 nucleotides and can have or not a limited protein-coding capability (Ulitsky *et al.*, 2013). They are limited to some specific cell types, playing crucial roles in tumorigenesis, because they modulate key pathways in transcriptional, posttranscriptional and epigenetic levels (Quinn *et al.*, 2016).

Due to their tissue specificity, they could potentially be used to distinguish types of malignancies (Flippot *et al.*, 2016).

Some lncRNAs are already used as diagnostic biomarkers in prostate, hepatocellular (Yuan *et al.*, 2017), colorectal (Zhao *et al.*, 2015) and lung cancer (Ren *et al.*, 2013). Regarding pancreatic cancer specifically, lncRNAs are diversely expressed according to tumor stage (primary or metastatic cancer) (Tahira *et al.*, 2011).

For early-stage diagnosis, some lncRNAs that are dysregulated in pancreatic cancer comparing with healthy pancreatic tissues could be potentially used as biomarkers as are the cases of H19 (Scaiewicz *et al.*, 2010), HOTAIR (Kim *et al.*, 2013), HOTTIP (Fu

*et al.*, 2017), MALAT-1 (Jiao *et al.*, 2014), HULC (Peng *et al.*, 2014), and GAS5 (Lu *et al.*, 2013).

Fragments of the lncRNAs HOTTIP-005 and RP11-567G11.1, namely HDRF and RDRF should also be potential biomarkers, either each alone or in combination (Wang *et al.*, 2015).

Most recently, Giulietti *et al.* (2018) identified eleven lncRNAs (A2M-AS1, DLEU2, LINC01133, LINC00675, MIR155HG, SLC25A25-AS1, LINC01857, LOC642852 (LINC00205), ITGB2-AS1, TSPOAP1-AS1 and PSMB8-AS1) that have different expression and promoter methylation levels and copy number alteration when comparing pancreatic cancer patients with healthy controls. Some of these, namely A2M-AS1, LINC01133, LINC00205 and TSPOAP1-AS1, could also be used as prognostic biomarkers.

Continuing to study lncRNAs may help understand the molecular pathogenesis of pancreatic cancer (table 10).

Table 10. lncRNA with potential as biomarkers

Biomarker Panel (concentration)	AUC, sensitivity, specificity	Reference
<b>H19</b>	Not validated yet	Scalewicz et al., 2010
<b>HOTAIR</b>	Not validated yet	Kim et al., 2013
<b>HOTTIP</b>	Not validated yet	Fu et al., 2017
<b>MALAT-1</b>	Not validated yet	Jiao et al., 2014
<b>HULC</b>	Not validated yet	Peng et al., 2014
<b>GAS5</b>	Not validated yet	Lu et al., 2013
<b>HDRF and RDRF</b>	Not validated yet	Wang et al., 2015
<b>A2M-AS1, DLEU2, LINC01133, LINC00675, MIR155HG, SLC25A25- AS1, LINC01857, LOC642852 (LINC00205), ITGB2-AS1, TSPOAP1- AS1 and PSMB8-AS1</b>	Not validated yet	Giulietti et al., 2018

### 3.1.6 Inflammatory factors and growth factors

It is a known fact that inflammation plays an important role in tumorigenesis and metastasis that is why inflammatory factors like chemokines, cytokines and growth factors could be potential diagnostic biomarkers (Kaur *et al.*, 2012).

Due to the fact that to the development of a tumor there needs to occur a growth factor stimulation and neovascularization (Zhang *et al.*, 2018b), pancreatic cancer patients have high serum levels of two growth factors, namely VEGF and bFGF (Pistol-Tanase *et al.*, 2008). These high levels also correlate to tumor size.

Shaw *et al.* (2014), studied IP-10, IL-6, PDGF and CA19-9 improving its diagnostic performance.

After comparing several inflammatory factor profiles between healthy and pancreatic cancer patients, Wingren *et al.* (2012) found that several ones were overexpressed: C1 esterase inhibitor, C3, C5, CD40, CD40 ligand, factor B, GLP-1, IFN- $\gamma$ , igm IL-10, IL-11, IL-12, IL-13, IL-16, IL-18, IL-1-ra, IL-1 $\alpha$ , IL-3, IL-5, IL-6, IL-7 and IL-8, integrin-

$\alpha$ -11, procathepsin W, sialyl Lewis x, TGF- $\beta$ 1, TNF- $\alpha$  and VEGF, based on which he constructed a panel of 25 biomarkers for early-stage diagnosis (AUC 0.95).

Yoshinaga *et al.* (2018), researched angiopoietin-like protein 2 (ANGPTL2), which is related to chronic inflammation and Type 2 diabetes mellitus, regarding its capability to early-diagnosis. The serum concentration of ANGPTL2 was significantly higher in pancreatic cancer patients than in healthy controls making it a potential biomarker to be furtherly studied.

Macrophage Inhibitory Cytokine 1 (MUC-1) seems promising, showing high levels in pancreatic adenocarcinoma (O'Brien *et al.*, 2015). It also proved to have a good diagnostic accuracy when combined with CA19-9 (Koopmann *et al.*, 2004, Ni *et al.*, 2005) (table 11).

Table 11. Summary of inflammatory and growth factors

Biomarker Panel (concentration)	AUC, sensitivity, specificity	Reference
<b>IP-10, IL-6, PDGF and CA19-9</b>	Not validated yet	Shaw et al., 2014
<b>C1 esterase inhibitor, C3, C5, CD40, CD40 ligand, factor B, GLP-1, IFN-<math>\gamma</math>, igm IL-10, IL-11, IL-12, IL-13, IL-16, IL-18, IL-1-ra, IL-1<math>\alpha</math>, IL-3, IL-5, IL-6, IL-7 and IL-8, integrin-<math>\alpha</math>-11, procathepsin W, sialyl Lewis x, TGF-<math>\beta</math>1, TNF-<math>\alpha</math> and VEGF</b>	AUC: 0.95	Wingren et al., 2012
<b>ANGPTL2</b>	Not validated yet	Yoshinaga et al., 2018
<b>MUC-1</b>	Not validated yet	O'Brien et al., 2015
<b>MUC-1, CA19-9</b>	Not validated yet	Koopmann et al., 2004, Ni et al., 2005

### 3.1.7 Circulating tumor cells (CTCs)

Circulating tumor cells are tumor cells that can enter circulation (Herrerros-Vilanueva *et al.*, 2016). They can be detected in early stages (before imaging techniques (Rhim *et al.*, 2014)), in bloodstream as they leave primary lesion in the beginning of tumor development (Pantel *et al.*, 2004, Husemann *et al.*, 2008, Rhim *et al.*, 2012). Their existence is already known for a century (Riva *et al.*, 2016). Due to the fact that they carry tumor markers on the surface and somatic mutations, they can be used for liquid and quick biopsies (O'Flaherty *et al.*, 2012).

CTCs can be identified in at least 40% (with some studies suggesting that it can reach even 100% of patients with pancreatic cancer) (Iwanicki-Caron *et al.*, 2013).

They began to be studied only recently for the reason that the technological advances needed to its analysis were only made in the last times. CellSearch platform enabled the selecting of CTCs from blood cells, as they express cell adhesion molecule (EpCAM) (Nagrath *et al.*, 2007).

It is believed that they are responsible for metastasis development (Herrerros-Vilanueva *et al.*, 2016). Nevertheless, its isolation and enrichment are extremely difficult making its detection extremely rare in pancreatic cancer (Rhim *et al.*, 2012).

The results of the studies of CTCs still remain debatable. According to a study by Albuquerque *et al.* (Albuquerque *et al.*, 2012), 47% of pancreatic cancer patients presented CTCs. In contrast to this, another study (Bidard *et al.*, 2013), demonstrated that there were detected by CellSearch platform CTCs in only 5% of patients.

Yang *et al.*, (Gao *et al.*, 2016), analyzed CTCs with another technique, consisting in a platform with subtraction, enrichment and immunostaining-fluorescence *in situ* hybridization. It showed an 88% sensitivity, 90% specificity and an early-detection rate of 12/13.

It can therefore be concluded that CTCs are promising biomarkers, even if only as prognostic biomarkers (Riva *et al.*, 2016). There must still be developed a standardized detection method and a large-scale validation (Zhang *et al.*, 2018b).

### 3.1.8 Exosomes

Exosomes are small vesicles containing nucleic acids and proteins (Skog *et al.*, 2008, Trjkovic *et al.*, 2008). They are secreted by almost all cells, including cancer cells, playing an important role in intercellular communication, tumorigenesis and metastasis (Melo *et al.*, 2014, Robbins *et al.*, 2014 Hoshino *et al.*, 2015).

They are composed externally by a lipid bilayer and do not have any cellular organelles (Azmi *et al.*, 2013). When released, they are very stable in the extracellular environment and can also be taken up by other cells, exchange material or information between cells (Lo Cicero *et al.*, 2015), and promote tumorigenesis (Charrier *et al.*, 2014). The main circulating DNA is associated with exosomes (Kahlert *et al.*, 2014). Exosomes contain many proteins, nucleic acids and lipids from cancer cells making them good candidates for diagnostic biomarkers. They are stable in almost all body fluids. What makes them good candidates as biomarkers is that they enter circulation in early stages of the tumorigenesis. In the case of pancreatic cancer, metastasis can even occur at early stage (Zhang *et al.*, 2018b). However, the isolation of malignant exosomes is challenging, making the development/improvement of techniques crucial (table 12).

#### **3.1.8.1 Proteins**

Glypican-1 (GPC1) is a membrane-anchored protein that is overexpressed in pancreatic cancer and precancerous lesions. For differentiating healthy controls, it had a perfect AUC (1.0) (Melo *et al.*, 2015). Comparing sensitivities GPC1 from serum exosomes has a better sensitivity than from whole serum. Meanwhile there have already been enough studies proving its significance for diagnosing pancreatic cancer (Zoller *et al.*, 2013).

Combining five proteins (CD44v6, Tspan8, epcam, MET, and CD104) and four miRNAs (miR-1246, miR- 4644, miR-3976, and miR-4306) in circulating exosomes it could discriminate pancreatic cancer with a sensitivity of 1.0 and a specificity of 0.8 (Madhavan *et al.*, 2015).

#### **3.1.8.2 DNA with somatic mutations**

Circulating exosomes carry a large amount of tumor DNA with its variations, somatic mutations and expressed fusion genes (San Lucas *et al.*, 2016).

As already approached, during malignancies, there occur and are identifiable KRAS and TP53 mutations. These are also present in circulating exosomal DNA, with a detection rate of almost 40% for pancreatic cancer patients, almost 30% in precancerous lesions and only 3% in healthy controls (Yang *et al.*, 2017).

The detection of KRAS mutations is higher in exosomes than in cfDNA (Allenson *et al.*, 2017).

### 3.1.8.3 miRNAs

Ding *et al.* (2015) disclosed that exosomes from pancreatic cancer transfer miRNAs to dendritic cells by miR-212 to induce immunotolerance.

Regarding the detection miRNA in circulating exosomes, pancreatic cancer patients have different levels of miR-10b, miR-20a, miR-21, miR-30c, miR-106b and miR-let7a. These levels normalized after tumor resection (Lai *et al.*, 2017).

Table 12. Principal exosomes

Biomarker Panel (concentration)	AUC, sensitivity, specificity	Reference
Proteins		
GPC1	AUC: 1.0	Melo <i>et al.</i> , 2015
CD44v6, Tspan8, epcam, MET, CD104, miR-1246, miR- 4644, miR-3976, miR-4306	Sensitivity: 100% Specificity: 80%	Madhaven <i>et al.</i> , 2015
DNA with somatic mutations		
TP53	Detection rate: 40% of pancreatic cancer patients	Yang <i>et al.</i> , 2017
KRAS		
miRNAs		
miR-10b, miR-20a, miR-21, miR-30c, miR-106b and miR-let7a	Aberrant levels	Lai <i>et al.</i> , 2017

### 3.1.9 Soluble stroma-related Biomarkers

Stromal modifications occur early in tumorigenesis and persist. Tumor initiation, growth and metastasis relies on the creation of a favorable microenvironment, including cancer cells, stroma and immune/ inflammatory cells (Erkan *et al.*, 2013), reason why, stroma-related circulating molecules make good potential early-stage biomarkers.

Resovi *et al.* (2018), studied 38 potential molecules (extracellular matrix proteins and proteolytic fragments, matrix-degrading enzymes and their inhibitors, growth factors, antiangiogenic factors, adhesion molecules (Yu *et al.*, 2005; Bloomston *et al.*, 2006;



Faca *et al.*, 2008; Kojima *et al.*, 2008; Fiedler *et al.*, 2009; Rong *et al.*, 2010; Xue *et al.*, 2010; Pan *et al.*, 2011)).

Of these, seven molecules were significantly up-regulated in pancreatic cancer when compared with healthy controls, namely MMP7 (AUC 0.98) with an excellent discriminatory ability, CCN2 (AUC 0.86) similar to CA19-9 (AUC 0.87), TIMP1 (AUC 0.82), IGFBP2 (Insulin Like Growth Factor Binding Protein 2) (AUC 0.82), TSP2 (Thrombospondin-2) (AUC 0.78), sICAM1 (Soluble intercellular adhesion molecule-1) (AUC 0.77) and PLG (plasminogen) (AUC 0.66).

Resovi *et al.* (2018), also studied combination of these seven potential biomarkers, concluding that the panel consisting of MMP7, CA19-9 and CCN2 had the best results (AUC 0.94).

MMP7 (matrilysin) is an inducer of acinar-to-ductal metaplasia (Crawford *et al.*, 2002; Sawey *et al.*, 2007), that is upregulated in precancerous (pancreatic) lesions. There is known to be high MMP7 plasma levels in association with early tumor progression in precancerous and early-stages (I, II), reason why it has value as a potential early diagnostic biomarker.

CCN2 (connective tissue growth factor (CTGF)) is overexpressed by activated pancreatic stellate cells in the earliest stages of pancreatic cancer. It promotes local desmoplasia, tumor survival, and metastasis (Leask *et al.*, 2009) (table 13).

Table 13. Soluble stroma-related biomarkers (summary)

Biomarker Panel (concentration)	AUC, sensitivity, specificity	Reference
<b>MMP7</b>	AUC 0.98	Resovi et al., 2018
<b>CCN2</b>	AUC 0.86	Resovi et al., 2018
<b>TIMP1</b>	AUC 0.82	Resovi et al., 2018
<b>IGFBP2</b>	AUC 0.82	Resovi et al., 2018
<b>TSP2</b>	AUC 0.78	Resovi et al., 2018
<b>siCAM1</b>	AUC 0.77	Resovi et al., 2018
<b>PLG</b>	AUC 0.66	Resovi et al., 2018
<b>MMP7, CA19-9 and CCN2</b>	AUC 0.94	Resovi et al., 2018

### 3.2 Glycosylation in Cancer

Glycosylation is the enzymatic process in which saccharides are linked through a glycosidic linkage to other saccharides, proteins or lipids (Fuster *et al.*, 2005, Moremen *et al.*, 2012). Glycoproteins consist on one or more glycans covalently attached to a polypeptide backbone. This linkage usually occurs via nitrogen or oxygen linkages, reason why they are named *N*-glycans or *O*-glycans, respectively (Varki *et al.*, 2009, Bennett *et al.*, 2012).

Glycan structures of glycoconjugates such as glycoproteins are modified in diseases, like inflammation, infectious diseases, diabetes, neurodegeneration and cancer (Kuzmanov *et al.*, 2013, Hart *et al.*, 2015, Munkley *et al.*, 2016, Kailemia *et al.*, 2017). Alterations in glycosylation in cancer have already been described for over 60 years (Ladenson *et al.*, 1949, Hakomori *et al.*, 1968). One of the main features in tumorigenesis seem to be aberrant glycosylation, as glycans are involved in main regulatory mechanisms, like protein folding and clearance rates, cell signaling (Boscher *et al.*, 2011, de-Freitas-Junior *et al.*, 2013, Gomes *et al.*, 2013, Takeuchi *et al.*, 2014), angiogenesis, differentiation, cell growth, cell-matrix interactions (Zhao *et al.*, 2008), immune modulation (Zhao *et al.*, 2008), tumor cell dissociation, invasion,

epithelial-mesenchymal transition (EMT), and metastasis (Pinho *et al.*, 2015, Llop *et al.*, 2018). Glycans alter protein conformation and structure, modulating in this way the functional activity of the protein (Helenius *et al.*, 2001).

For a tumor to develop it has to gain the ability to overcome cell–cell adhesion and to be able to invade surrounding tissue. Epithelial cadherin (E-cadherin) is a transmembrane glycoprotein (Pinho *et al.*, 2011) and a major epithelial cell–cell adhesion molecule in cancer (Paredes *et al.*, 2012). Glycans can have a crucial effect on tumor cell–cell adhesion by directly interfering with E-cadherin functions. In cancer there is also a mutual regulatory mechanism between E-cadherin-mediated cell–cell adhesion and its glycosylation leading either to tumor suppression or tumor metastasis (Gu *et al.*, 2009, Pinho *et al.*, 2013).

Cancer cells have a different metabolism than normal ones. There is a shift from oxidative phosphorylation to aerobic glycolysis, called the Warburg effect (Warburg *et al.*, 1956). This high glucose uptake is used to cope with the increased energetic and biosynthetic needs to generate a tumor. For this purpose, cancer cell can also upregulate glutamine uptake. As a result, the cytoplasm of cancer cells is abundant in glucose. It contributes to increased glycolysis and increases the flux into the metabolic branch pathways.

### **3.2.1 Glycomic aberration vs. biomarkers**

As alterations in glycosylation regulate the development and consequent progression of cancer, they are potential biomarkers for early-diagnosis and targets for therapeutic strategies (Hakomori *et al.*, 2002, Fuster *et al.*, 2005, Taniguchi *et al.*, 2009, Reis *et al.*, 2010, Freeze *et al.*, 2013, Pinho *et al.*, 2013). Alterations in glycosylation have a major heterogeneity. This is because aberrant glycan modifications are protein-specific, site-specific (meaning that different sites on the protein can be differentially glycosylated) and cell-specific (Pinho *et al.*, 2015). There are two main mechanisms for tumor-glycosylation firstly postulated by Hakomori and *et al.* (1983), namely incomplete synthesis and neo-synthesis process. Regarding incomplete synthesis process, it occurs normally in early-stages of the malignancy. It is a consequence of the damage of the normal synthesis of complex glycans and leads to the biosynthesis of truncated structures (abnormally shortened glycans), like sialyl Tn. In contrast to this, neo-synthesis process more often arises in advanced stages. It regards the induction of genes involved in expression of carbohydrate determinants (like *de novo* expression of antigens like sialyl Lewis). These modifications can be detected in biological fluids of the cancer patients, as a result of the aberrant glycoproteins being

shed into the bloodstream (Peracaula *et al.*, 2008, Meany *et al.*, 2011). All of the nine proteins (Polasinski *et al.*, 2007) approved as cancer biomarkers, are glycosylated proteins (Schiess *et al.*, 2009, Diamandis *et al.*, 2010, Füzéry *et al.*, 2013, Kuzmanov *et al.*, 2013, Pavlou *et al.*, 2013) showing the importance of them.

There is a major glycan diversity, due to many factors (Pinho *et al.*, 2015). First of all, monosaccharides can have innumerable compositions, as is the case of galactose or *N*-acetylgalactosamine for example. Monosaccharides can also be linked in different ways, with the Carbon atoms involved in the linkage varying (C1-C3 or C1-C4). In addition to that, monosaccharides have a terminal aldehyde/ketone. This hydroxy group can either be in  $\alpha$  or  $\beta$  configuration, changing accordingly the anomeric state of the monosaccharide. Branching structures, or in its linkage to their aglycone (non-glycosyl) part (Cummings *et al.*, 2009, Varki *et al.*, 2009). Concerning abnormal glycosylation, this too has innumerable causes. In the first place it can be due to under- or overexpression of glycosyltransferases (enzymes that catalyze the transfer of the glycosyl-group to the aglycone (lipids, proteins, carbohydrates)). This occurs because of a dysregulation at transcriptional level (Kannagi *et al.*, 2009 Hatano *et al.*, 2011, Pinho *et al.*, 2012), a dysregulation of chaperone function (Schietinger *et al.*, 2006, Aryal *et al.*, 2010) (chaperones stabilize unfolded proteins) or altered glycosidase activity (Kakugawa *et al.*, 2002) (enzymes that catalyze hydrolysis of glycosylic linkages, separate glycosyl-group from aglycone). Furthermore, abnormal glycosylation can be caused by changes in tertiary conformation of the peptide backbone and of the nascent glycan chain. Another possible cause consists on the variability of acceptor substrates in addition to the availability and abundance of sugar nucleotide donors and cofactors (Kumamoto *et al.*, 2001). This means that there is an alteration in glycan expression, because there is a whole panoply of ligands that can be linked during glycosylation. Besides, glycosyltransferases can be mislocated (in the Golgi apparatus) or there can be a change in its activity leading to synthesis of immature core glycan structures (Kellokumpu *et al.*, 2001, Marcos *et al.*, 2004, Brockhausen *et al.*, 2006, Gill *et al.*, 2010).

There are many studies regarding serum glycoproteins with altered glycan chains in pancreatic cancer patients, especially altered N-glycan patterns in tumors (Arnold *et al.*, 2008). They do not have sufficient specificity though, as many of these N-glycosylation also occur in benign conditions. Llop *et al.* (2018) propose combining aberrant glycosylation with the altered protein levels in order to improve its performance in diagnosing pancreatic cancer (Kailemia *et al.*, 2018).

Almost all tumor cells tend to be affected by aberrant glycosylation, making these changes in glycosylation more pronounced than alterations in protein expression.

With progression of the cancer, glycans expressed are continuously and rapidly changing (Pinho *et al.*, 2015, Munkley *et al.*, 2016) and alterations become more marked the more aggressive the tumor becomes. This is why glycan alterations should be more reliable biomarkers, also in predictive value (Silva *et al.*, 2015).

Nonetheless, the analysis of the glycoproteome in pancreatic cancer has not been extensively studied yet. One possible reason for this is that the analysis of the glycoproteome is much more complex than the one of the proteome, as there is no template (like the amino acid sequence coded in the genome for proteins) for glycans (Llop *et al.*, 2018) and there is a major amount of heterogeneity.

There are four main cancer-associated glycosylation processes (Stowell *et al.*, 2015): sialylation (sialyl-Lewis<sup>x</sup>= CA19-9 antigen), fucosylation, increased GlcNAc-branching of N-glycans and over-expression of truncated mucin-type O-glycans.

Sialylation plays a critical role in cellular recognition, cell adhesion and signaling. There are two major sialylated antigens associated with cancer, namely SLe<sup>a</sup>, SLe<sup>x</sup>. Both of them are correlated with a poor prognosis (Amado *et al.*, 1998, Baldus *et al.*, 1998).

SLe<sup>x</sup> is a ligand for selectins (Rosen *et al.*, 1994). Selectins are vascular cell adhesion and belong to a family of C-type lectins. During inflammation selectins mediate the initial attachment of leukocytes to the endothelium during the process of leukocyte extravasation (Rosen *et al.*, 1994). In cancer, SLe<sup>x</sup> interactions with selectins regulate the metastatic cascade, determining the malignant behavior and development of metastasis (Nakamori *et al.*, 1993).

Ceruplasmin is an acute-phase protein, that has an increased sialyl-Lewis (SLe<sup>x</sup>) level in pancreatic cancer. In a big cohort study of pancreatic cancer serum samples, the ratio of SLe<sup>x</sup> on ceruplasmin related to the ceruplasmin level showed to be increased in pancreatic cancer in comparison to chronic pancreatitis (Balmaña *et al.*, 2015). SLe<sup>a</sup> is detected by serological assay CA19-9.

Concerning fucosylation, it is catalyzed by numerous fucosyltransferases, including FUT3, which is a Lewis Gene (Le). Fucosylation is divided in terminal and core fucosylation. Terminal fucosylation originates specific Lewis blood-group antigens (Le<sup>x</sup> and Le<sup>y</sup> and Le<sup>a</sup> and Le<sup>b</sup>). Contrarily, core fucosylation consists in the addition of  $\alpha$ 1,6-fucose to the innermost GlcNAc residue of N-glycans through the action of Fuc-TVIII (encoded by *FUT8*) (Carvalho *et al.*, 2010).

Otherwise, branching and bisecting GlcNAc N-glycans is also commonly present in malignancies. In these cases, there is an increase in the expression of complex  $\beta$ 1,6-branched N-linked glycans 2 (Dennis *et al.*, 1987). Branched N-glycans are further modified by  $\beta$ 1,4-GalTs, elongated with poly-N-acetylactosamine and further capped

with sialic acid and fucose (Dennis *et al.*, 1987). This poly-*N*-acetylactosamine structure is a ligand for galectins (Di Lella *et al.*, 2011). Galectins have important roles in cancer, contributing to neoplastic transformation, tumor cell survival, angiogenesis and tumor metastasis (Croci *et al.*, 2014).

$\alpha$ -fetoprotein (AFP) is a glycoprotein fucosylated and in overexpression in hepatocellular carcinoma and other benign hepatic diseases. Detecting simultaneously both its overexpression and its fucosylation (ratio fucosylated AFP/ total AFP) is approved as a biomarker for risk assessment of risk patients for the development of hepatocellular carcinoma (Li *et al.*, 2001, Wang *et al.*, 2009). With the same thought, there exists also biomarker with changes in fucosylation for prostate cancer (Tabarés *et al.*, 2006, Sarrats *et al.*, 2010) used for screening high-risk populations (Llop *et al.*, 2016, Ferrer-Batallé *et al.*, 2017). Both biomarkers already used in two types of cancer show the great potential of fucosylation in early-stage diagnostics of cancer.

Regarding the glycosylation profile of pancreatic carcinomas, there is an increase of branching of N-linked oligosaccharides (Zhao *et al.*, 2007), in protein fucosylation and sialylation in pancreatic cancer serum. Another main feature consists on the increment in Lewis and blood group glycans (Pour *et al.*, 1988, Pérez-Garay *et al.*, 2013). According to Remmers *et al.* (2013), there also is also present a truncated O-linked glycosylation resulting in the Tn and sialyl-Tn antigens.

As a starting point for the search of glycoproteins as biomarkers normally count whole serum samples or serum after removing the most abundant ones are removed, as they should hinder the detection aberrant glycosylation patterns that should exist in low concentration (Llorens *et al.*, 2018). There can be used lectins or antibodies against specific glycan structures or mass-spectrometry (MS) to identify the different glycosylation of the glycoproteins.

Normally the strategy consists on analyzing alterations in branching, fucosylation (Tan *et al.*, 2015, Terao *et al.*, 2015) sialylation (Kontro *et al.*, 2014) and specific N-glycosylation occupancy (Pan *et al.*, 2014).

There is an increase in fucosylation and sialylation described for pancreatic cancer (Zhao *et al.*, 2007, Li *et al.*, 2009).

Two acute-phase proteins (haptoglobin (HPT) and  $\alpha$ -1-acid-glycoprotein (AGP)) are increased in core fucosylation in pancreatic cancer, when comparing to healthy controls and chronic pancreatitis (Sarrats *et al.*, 2010). However, according to (Matsumoto *et al.*, 2010), the increase in fucosylated HPT has no significant difference from chronic pancreatitis patients. However, combining them with CA19-9 could improve the diagnostic potential in contrast to the both of the biomarkers isolated. AGP

was also found to be increased in its  $\alpha$ -1,3-fucose glycoforms (Giménez *et al.*, 2015, Balmaña *et al.*, 2016, Mancera-Arteu *et al.*, 2017).

$\alpha$ 1-Antichymotrypsin (ACT), trombospondin-1 and HPT were the best performing fucosylated glycoproteins in a study by Nie *et al.* (2014) in which lectins (*Aleuria aurantia lectin*) were used to analyze the samples. Their combination with CA19-9 proved to have a high diagnostic potential.

ACT also showed a good potential, when analyzed by MS in another study (Tan *et al.*, 2015).

Although isolated levels of serum pancreatic RNase 1 are not enough to consider it a potential biomarker, this protein is much more core fucosylated in pancreatic cancer. Quantification of its core fucosylation could therefore be a good biomarker (Barrabés *et al.*, 2007). As it was shown that the Asn-88 occupation site is significantly increased in N-glycosylation in pancreatic cancer, this Asn-88 site could also be a novel biomarker (Nakata *et al.*, 2014).

Regarding now N-glycan branching of serum glycoproteins, Drabik *et al.* (2017) found four glycoproteins (LIFR, CE350, VP13A and HPT) that carrying an altered N-glycan structure in pancreatic cancer.

Mucins can be secreted from tissues to the bloodstream, making its specific glycoforms easy to detect in serum. Altered glycosylation patterns of mucin (MUC), especially MUC4 (glycoform found in neoplasms: MUC4-Tn) and MUC1 (glycoform found in neoplasms: MUC1-STn) have a good potential as biomarkers (Remmers *et al.*, 2013). Immunoprecipitated MUC1 can be studied using lectins (Matsuda *et al.*, 2017) or antibody-lectin sandwich arrays with which it was found to carry the CA19-9 antigen (SLe<sup>a</sup>) in pancreatic cancer patients (Yue *et al.*, 2009). Alterations on MUC1 and CEA were also found by Chen *et al.* (Chen *et al.*, 2007) using antibody microarrays.

MUC5AC present in cyst fluid samples can be detected with wheat germ agglutinin lectin and according to Haab *et al.* (2010), can identify malignant pre-cancerous stages.

All of these mucins were described in pancreatic cancer in tissue and not serum (that is the most convenient sample). Due to analytical limitations, O-glycans in mucins cannot be analyzed yet (Llop *et al.*, 2018).

Recently reported was also an increased expression of MUC1-SLe<sup>x</sup> and MUC5AC-SLe<sup>x</sup> (Balmaña *et al.*, 2018). MUC5AC-SLe<sup>x</sup> is observed in serum from pancreatic cancer patients carrying 3 fucose (Singh *et al.*, 2015). Its performance is improved when in comparison with CA19-9 (Tang *et al.*, 2016).

Concluding, most of the studies focused on finding altered glycosylation in serum or depleted serum samples. It has mostly resulted in the identification of acute-phase glycoproteins and mucins (MUC1 and MUC5AC). The changes of these, occurred in sialic acid, fucose of SLe<sup>x/a</sup> occurring mostly in advanced stages of the malignancy or inflammatory stages, limiting its use as early-stage biomarkers. However, combining these aberrant glycolisation with protein levels and imaging could improve its performance (table 14).



Table 14. Glycosylation Biomarkers

Biomarker Panel	Characteristics	Reference
<b>HPT, AGP, CA19-9</b>	Increased core fucosylation	Sarrats <i>et al.</i> , 2010 Matsumoto <i>et al.</i> , 2010
<b>HPT, ACT, CA19-9</b>	Increased core fucosylation	Nie <i>et al.</i> , 2014 Tan <i>et al.</i> , 2015
<b>RNase 1</b>	Increased core fucosylation	Barrabés <i>et al.</i> , 2007
<b>Asn-88 occupation site</b>	Increased in N-glycosylation	Nakata <i>et al.</i> , 2014
<b>LIFR, CE350, VP13A and HPT</b>	Altered N-glycan branching	Drabik <i>et al.</i> , 2017
<b>MUC4 (glycoform found in neoplasms: MUC4-Tn)</b>	Altered glycosylation patterns	Remmers <i>et al.</i> , 2013
<b>MUC1 (glycoform found in neoplasms: MUC1-STn)</b>	Altered glycosylation patterns	Remmers <i>et al.</i> , 2013 Chen <i>et al.</i> , 2007
<b>MUC5AC</b>	Altered glycosylation patterns	Haab <i>et al.</i> , 2010
<b>MUC1-SLe<sup>x</sup></b>	Increased expression	Balmaña <i>et al.</i> , 2018
<b>MUC5AC-SLe<sup>x</sup></b>	Increased expression, carrying 3'fucose	Balmaña <i>et al.</i> , 2018 Singh <i>et al.</i> , 2015
<b>MUC1 and MUC5AC, protein levels and imaging</b>	Changes in sialic acid, fucose of SLe <sup>x/a</sup> in advanced stages hence the combination	Llop <i>et al.</i> , 2018

### 3.3 Lectins as a tool

Lectins are ubiquitous proteins. They show a specificity for the carbohydrate moiety of glycoconjugates (Hashim *et al.*, 2017) being defined as glycan-specific receptors. They are the so-called decoding system for the oligosaccharide codes (Ribeiro *et al.*, 2018). It has been difficult to classify lectins, due to their immense diversity. However, nowadays lectins are classified according to distinct protein folding, domains/structural similarities and evolutionary-relatedness of proteins (Peumans *et al.*, 2001). According to these criteria, there are twelve lectin families, (Van Damme *et al.*, 2008). The first lectin to be discovered is believed to have been ricin in 1888 (Sharon *et al.*, 2004). It

is an extremely toxic protein and can be found in *Ricinus communis*. Like ricin, most of lectins have a very high cytotoxicity. Since then, there are more than a thousand plants reported to possess lectins and it is believed that a large number still remains to be discovered (Ribeiro *et al.*, 2018). Due to their carbohydrate binding specificities, lectins have been thoroughly studied, having innumerable applications now-a-days. Lectins can be found in plants (mostly as seed storage proteins (Ribeiro *et al.*, 2014)) and fungi and very dispersed in nature.

### **3.3.1 Biomedical Application**

In biomedical research, lectins play an important role. Whether in cancer, as a means of discovering new biomarkers, whether in neurodegenerative diseases or as a means of target-therapy (Ribeiro *et al.*, 2018).

Due to the specificity of lectins, they can recognize carbohydrates on the cellular membrane, making them excellent tools for decoding the abnormal glycosylation happening in malignancies. Many of these aberrantly glycosylated expressed peptides in cancer patients, are not possibly detected using conventional methods due to the fact that they occur in very low concentrations. In combination with mass spectrometry, lectins can help identifying several potential cancer biomarkers, based on glycosylation degree (Mody *et al.*, 1995). Most of the methods in which lectins are used to identify aberrations in glycolysation in malignancies consist on adaptations of other methods normally used with antibodies (Enzyme-linked lectin assay, lectin histochemistry, lectin blotting).

Furthermore, it has also been found that there are some lectins with anticancer properties. As a result of lectins dictating cytotoxic effects and mediating apoptosis (Park *et al.*, 2000) and autophagy they can prevent tumor growth.

Programmed cell death can also be one effect caused by lectins, when used as anticancer therapy (Ribeiro *et al.*, 2018).

Lectins also have great potential as tissue specific therapy due to its specificity for a specific oligosaccharide side chain (Ribeiro *et al.*, 2018).

### **3.3.2 Immobilized-Lectin Affinity Chromatography**

Immobilized-lectin affinity chromatography consists on immobilizing lectins onto a matrix and its carbohydrate ligands (figure 12). It is a method making use of the specificity of lectins interactions aiming for glycoproteins separation and enrichment (Hage *et al.*, 2012). As a technique used for identifying potential biomarkers, it is

normally used in addition with mass spectrometry analysis. By running a sample, the result is a glycoprotein-enriched eluate. Comparing, a healthy control and a pancreatic cancer patient samples, there can easily be identified the aberrantly expressed or glycosylated glycoproteins. For pancreatic cancer, there has been identified a potential biomarker with this technique. The enrichment of core fucosylated glycoproteins was carried out using *Lens culinaris* agglutinin in combination with mass spectrometry analysis (Tan *et al.*, 2015).

Another used variant is the use of a panel of lectins, instead of just one. Making use of the complement of the panel multi-lectin affinity chromatography gives a deeper analysis (Hashim *et al.*, 2017).




Figure 12. Immobilized-lectin affinity chromatography

(adapted from Hashim *et al.*, 2017)

### 3.3.3 Enzyme-linked Lectin Assay

Enzyme-linked lectin assay consists in the replacement of antibodies by lectins in the enzyme-linked immunosorbent assay (Hashim *et al.*, 2017). The principle adopted remains the same. The enzyme-linked immunosorbent assay (McCoy *et al.*, 1983) is a direct assay in which to the sample in a microtiter plate is added an enzyme-conjugated lectin. While the lectin specifically binds to the glycan moiety, the enzyme converts the substrate into a colored product. The intensity of the coloration is measured with the help of a spectrophotometer and used to estimate the glycoconjugates level. This method is widely used, including in the research of cancer

biomarkers (Kuzmanov *et al.*, 2013), as it is quick, easy and requires only small amounts of samples. However, the correct identification of glycoproteins may only be possible by coupling enzyme-linked lectin assay with proteomics analysis or antibody detection.

For this method, there are three different and possible approaches (figure 13): (a) The direct assay, the classical protocol, as already told consists on the addition of an enzyme-conjugated lectin to the microtiter plate containing the sample. (b) In the hybrid assay it is the other way around, the plate contains the antibody capturing the specific glycoproteins, only after this is added the enzyme-conjugated lectin. (c) Sandwich enzyme-linked lectin assay involves two lectins and is almost a combination of the first two methods. One lectin is on the plate and captures the glycoprotein. The second lectin is used as a detection reagent.

Ching *et al.* (1989) identified a glycoprotein in the serum of pancreatic cancer patients that binds to peanut lectin. With the purpose of identifying it and diagnosing patients, Ching *et al.* (1989) developed a direct enzyme-linked PNA assay. The results (in terms of sensibility and specificity) were similar to the ones using CA19-9.




Figure 13. Enzyme-linked lectin assay approaches

(adapted from Hashim *et al.*, 2017)

### 3.3.4 Lectin Histochemistry

Lectin histochemistry is very similar to immunohistochemistry. It too consists in visualizing cellular components of tissue microscopically, but the antibodies are replaced by lectins (Hashim *et al.*, 2017). It can be used to obtain information about the glycosylation present in the sample. Comparing healthy controls with cancer samples may highlight aberrant glycosylations. There are two variants possible (figure 14): (a) the direct method, in which a lectin is covalently bonded to fluorophores, enzymes, colloidal gold or ferritin. (b) in the indirect method, a hapten (alone cannot

induce an immune response but when linked to a higher molecular weight carrier protein generates one), namely a biotin or digoxigenin is conjugated with a lectin, they are then recognized by an enzyme linked-streptavidin (highly selective to biotin making one of the strongest links) or -anti-digoxigenin (Roth *et al.*, 2011).

Lectin histochemistry has already been widely used for studying glycolysation aberrations in malignancies (Sobral *et al.*, 2010).




Figure 14. Lectin histochemistry

(adapted from Hashim *et al.*, 2017)

### 3.3.5 Lectin Blotting

As a result of replacing the antibodies used in western blotting by lectins arises the lectin blotting (Shan *et al.*, 2001). As in the western blot, samples are placed on a polyacrylamide gel and transferred (using current) onto a polyvinylidene fluoride or nitrocellulose membrane (figure 15). Glycoproteins are then detected by lectins and the visualization of the complex is possible by the use of conjugates (enzymes, biotin, digoxigenin, ...), like in histochemistry. Lectin blotting has been used for characterizing glycan structures (Akama *et al.*, 2006), detecting and quantifying N- or O-glycosylated proteins (Roth *et al.*, 2012) and detecting aberrations in glycosylation (like it happens in cancer) (Kitamura *et al.*, 2003). However, it does not have very potential in routine diagnostics (Hashim *et al.*, 2017).




Figure 15. Lectin blotting

(adapted from Hashim *et al.*, 2017)

### 3.3.6 Lectin Array

On the lectin array, multiple lectins are immobilized onto a solid support. Due to the fact that there are different lectins, with different specificities, different glycoproteins can be detected at the same time (Hu *et al.*, 2009, Hirabayashi *et al.*, 2011). It is a quick and sensitive analysis of glycans (Hashim *et al.*, 2017). Multiple potential biomarkers have already been identified with the help of lectin array, namely *Agaricus bisporus* lectin for colorectal cancer (Nakajima *et al.*, 2015). Lasectin array has been modified several times already. To this point, there are two variants being used (figure16): (a) one lectin per spot are organized into the lectin array slide. When the samples come in contact with the lectins, the spots where the specific glycoproteins are illuminated under appropriate scanner due to their interactions. (b) another

possibility is to conjugate lectins to different fluorescent colored beads. After entering in contact with samples, the beads pass through a detector with two lasers, one identifies the beads (classification laser) and the other quantifies them (reporter laser) (Hashim *et al.*, 2017).




Figure 16. Lectin array

(adapted from Hashim *et al.*, 2017)

## 4 Conclusions

Pancreatic cancer remains an extremely challenging disease. It is asymptomatic, lacks radiological manifestations and there still is no known specific molecule circulating in body fluids. It is only curable in early stages, showing no symptoms by this time.

To distinguish the malignancy in early-stage with adequate sensitivity and specificity, screening first with biomarkers and then imaging could be the solution. Development of a screening protocol for high-risk populations is crucial to enhance life expectancy. However, there are innumerable possible biomarkers.

As problems in development of biomarkers there is the fact that most patients are diagnosed in advanced stages. Due to this, having samples from early-stage available for screening for biomarkers is even more difficult. A possible way to overcome this,

could be to screen high-risk populations with positive family history. Controlling their biomarker levels once a year could be a possibility to detect a significant and specific increase (or decrease) in biomarkers, which could be used in clinical practice as early-stage biomarkers.

To evaluate its validity, these studies also need controls, specifically samples from benign pancreatic diseases. This could also not be easy to find.

The solution seems to come from developing a panel with CA19-9 and other biomarkers in order to improve sensitivity and specificity. CA19-9 is the only biomarker used in pancreatic cancer (for monitoring patients). Its main problem as an early-stage biomarker is due to the fact that it is not expressed in 10-20% of the Lewis antigen-negative population. Combining it in a panel with other biomarkers is promising. However, there still is a lot of work to be done in furtherly studying and validating the many different existing possibilities.

The most promising combinations can be seen table 15.



Table 15. Best performing potential biomarkers

<b>Biomarker Panel</b>	CA 19-9, albumin and IGF	CA 19-9, albumin, C-reactive protein and interleukin	CA19-9, TIMP1, LRG1	CA19-9, THBS2	Palmitic acid	Ezrin	CA19-9, miR-145, miR-150, miR-223, miR-636	CA19-9, miR-26b, miR-34a, miR-122, miR-126, miR-145, miR-150, miR-223, miR-505, miR-636, miR-885.5p	CD44v6, Tspan8, epcam, MET, CD104, miR-1246, miR-4644, miR-3976, miR-4306
<b>Results</b>	Sensitivity: 93,6%; Specificity: 95%	Sensitivity: 99.39%; Specificity: 90%	AUC: 0.95; Sensitivity: 75%; Specificity: 95%	Sensitivity: 87%; Specificity: 98%	AUC: 1.0; Sensitivity: 100%; Specificity: 100%	AUC: 0.9 Sensitivity: 93.2% Specificity: 75.5%	AUC: 0.93; Sensitivity: 85%; Specificity: 85%	AUC: 0.93; Sensitivity: 85%; Specificity: 85%	Sensitivity: 100% Specificity: 80%
<b>Reference</b>	Goh <i>et al.</i> , 2017	Zeh <i>et al.</i> , 2005	Capello <i>et al.</i> , 2017	Kim <i>et al.</i> , 2017	Di Gangi <i>et al.</i> , 2016	Capello <i>et al.</i> , 2013	Schultz <i>et al.</i> , 2014	Schultz <i>et al.</i> , 2014	Madhavan <i>et al.</i> , 2015

## **5 Challenges and Future Directions**

Finding an early-detection biomarker for pancreatic cancer passes by finding a combination of biomarkers that complement each other in order to achieve the best sensitivity and specificity. When using it in combination with imaging techniques, it should improve its results. Large-scale studies using the same standardized methods and conditions are needed to validate the results already obtained.

# Bibliography

Aberle DR, Adams AM, National Lung Screening Trial Research Team. (2011) Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med*; 365:395–409. doi: 10.1056/NEJMoa1102873.

Ahn SS, Kim MJ, Choi JY, Hong HS, Chung YE, Lim JS. (2009) Indicative findings of pancreatic cancer in prediagnostic CT. *Eur Radiol*; 19: 2448-2455.

Akama TO, Fukuda MN. (2006) N-Glycan structure analysis using lectins and an alpha-mannosidase activity assay. *Methods in Enzymology* 416:304–314. DOI 10.1016/s0076-6879(06)16020-6.

Al-Hawary MM, Francis IR, Chari ST, Fishman EK, Hough DM, Lu DS, Macari M, Megibow AJ, Miller FH, Morteale KJ, Merchant NB, Minter RM, Tamm EP, Sahani DV, Simeone DM. (2014) Pancreatic ductal adenocarcinoma radiology reporting template: consensus statement of the Society of Abdominal Radiology and the American Pancreatic Association. *Radiology*; 270: 248-260.

Albuquerque A, Kubisch I, Breier G, Stam-minger G, Fersis N, Eichler A, Kaul S and Stol-zel U. (2012) Multimarker gene analysis of circulating tumor cells in pancreatic cancer patients: a feasibility study. *Oncology*; 82: 3-10.

Allenson K, Castillo J, San Lucas FA, Scelo G, Kim DU, Bernard V, Davis G, Kumar T, Katz M, Overman MJ, Foretova L, Fabianova E, Holca- tova I, Janout V, Meric-Bernstam F, Gascoyne P, Wistuba I, Varadhachary G, Brennan P, Hanash S, Li D, Maitra A, Alvarez H. (2017) High prevalence of mutant KRAS in circulating exosome-derived DNA from early-stage pancreatic cancer patients. *Ann Oncol*; 28: 741-747.

Amado M, Carneiro F, Seixas M, Clausen H, Sobrinho-Simoes M. (1998) Dimeric sialyl-Le<sup>x</sup> expression in gastric carcinoma correlates with venous invasion and poor outcome. *Gastroenterology* 114, 462–470.

Arnold JN, Saldiva R, Hamid UM, Rudd PM. (2008) Evaluation of the serum N-linked glycome for the diagnosis of cancer and chronic inflammation. *Proteomics*; 8: 3284-3293. PMID: 18646009; DOI: 10.1002/pmic.200800163

Aryal RP, Ju T, Cummings RD. (2010) The endoplasmic reticulum chaperone Cosmc directly promotes in vitro folding of T-synthase. *J. Biol. Chem.* 285, 2456–2462.

Atkin WS, Edwards R, Kralj-Hans I, UK Flexible Sigmoidoscopy Trial Investigators. (2010) Once only flexible sigmoidoscopy screening in prevention of colorectal cancer: a multicentre randomised controlled trial. *Lancet*; 375:1624–1633.

Azmi AS, Bao B, Sarkar FH. (2013) Exosomes in cancer development, metastasis, and drug resistance: a comprehensive review. *Cancer Metastasis Rev*; 32: 623-642.

Baan R, Grosse Y, Straif K, Secretan B, El Ghissassi F, Bouvard V, Benbrahim-Tella L, Guha N, Freeman C, Galichet L, Coglian V, WHO International Agency for Research on Cancer Monograph Working Group. (2009) A review of human carcinogens-Part F: Chemical agents and related occupations. *Lancet Oncol*; 10: 1143-4.

Baldus SE, Zirbes TK, Mönig SP, Engel S, Monaca E, Rafiqpoor K, Hanisch FG, Hanski C, Thiele J, Pichlmaier H, Dienes HP. (1998) Histopathological Subtypes and Prognosis of Gastric Cancer Are Correlated with the Expression of Mucin-Associated Sialylated Antigens: Sialosyl-Lewis<sup>a</sup>, Sialosyl-Lewis<sup>x</sup> and Sialosyl-Tn. *Tumor Biol*;19:445-453. doi: 10.1159/000030036

Ballehaninna UK, Chamberlain RS. (2012) The clinical utility of serum CA 19-9 in the diagnosis, prognosis and management of pancreatic adenocarcinoma: An evidence-based appraisal. *J Gastrointest Oncol*; 3: 105-119. PMID: 22811878 DOI: 10.3978/j.issn.2078-6891.2011.021

Balmaña M, Duran A, Gomes C, Llop E, López-Martos R, Ortiz MR, Barrabés S, Reis CA, Peracaula R. (2018) Analysis of sialyl-Lewis x on MUC5AC and MUC1 mucins in pancreatic cancer tissues. *Int J Biol Macromol*; 112: 33-45. PMID: 29408556 DOI: 10.1016/j.ijbiomac.2018.01.148

Balmaña M, Giménez E, Puerta A, Llop E, Figueras J, Fort E, Sanz-Nebot V, de Bolós C, Rizzi A, Barrabés S, de Frutos M, Peracaula R. (2016) Increased  $\alpha$ 1-3 fucosylation of  $\alpha$ -1-acid glycoprotein (AGP) in pancreatic cancer. *J Proteomics*; 132: 144-154. PMID: 26563517 DOI: 10.1016/j.jprot.2015.11.006

Balmaña M, Sarrats A, Llop E, Barrabés S, Saldova R, Ferri MJ, Figueras J, Fort E, de Llorens R, Rudd PM, Peracaula R. (2015) Identification of potential pancreatic cancer serum markers: Increased sialyl-Lewis X on ceruloplasmin. *Clin Chim Acta*; 442: 56-62. PMID: 25595436 DOI: 10.1016/j.cca.2015.01.007

Barrabés S, Pagès-Pons L, Radcliffe CM, Tabarés G, Fort E, Royle L, Harvey DJ, Moenner M, Dwek RA, Rudd PM, De Llorens R, Peracaula R. (2007) Glycosylation of serum ribonuclease 1 indicates a major endothelial origin and reveals an increase in core fucosylation in pancreatic cancer. *Glycobiology*; 17: 388-400. PMID: 17229815 DOI: 10.1093/glycob/cwm002

Bauer AS, Keller A, Costello E, Greenhalf W, Bier M, Borries A, Beier M, Neoptolemos J, Büchler M, Werner J, Giese N, Hoheisel J. (2012) Diagnosis of pancreatic ductal adenocarcinoma and chronic pancreatitis by measurement of microRNA abundance in blood and tissue. *PLoS One*; 7: e34151.

<https://doi.org/10.1371/journal.pone.0034151>

Bennett, EP, Mandel U, Clausen H, Gerken TA, Fritz TA, Tabak LA. (2012) Control of mucin-type O-glycosylation: a classification of the polypeptide GalNAc-transferase gene family. *Glycobiology* 22, 736–756. doi: 10.1093/glycob/cwr182.

Bettegowda C, Sausen M, Leary RJ, Kinde I, Wang Y, Agrawal N, Bartlett BR, Wang H, Luber B, Alani RM, Antonarakis ES, Azad NS, Bardelli A, Brem H, Cameron JL, Lee CC, Fecher LA, Gallia GL, Gibbs P, Le D, Giuntoli RL, Goggins M, Hogarty MD, Holdhoff M, Hong SM, Jiao Y, Juhl HH, Kim JJ, Siravegna G, Laheru DA, Lauricella C, Lim M, Lipson EJ, Marie SK, Netto GJ, Oliner KS, Olivi A, Olsson L, Riggins GJ, Sartore-Bianchi A, Schmidt K, Shih IM, Oba-Shinjo SM, Siena S, Theodorescu D, Tie J, Harkins TT, Veronese S, Wang TL, Weingart JD, Wolfgang CL, Wood LD, Xing D, Hruban RH, Wu J, Allen PJ, Schmidt CM, Choti MA, Velculescu VE, Kinzler KW, Vogelstein B, Papadopoulos N, Diaz LA Jr. (2014) Detection of circulating tumor DNA in early- and late-stage human malignancies. *Sci Transl Med*; 6: 224ra24. PMID: 24553385 DOI: 10.1126/scitranslmed.3007094

Biankin AV, Waddell N, Kassahn KS, Gingras MC, Muthuswamy LB, Johns AL, Miller DK, Wilson PJ, Patch AM, Wu J, Chang DK, Cowley MJ, Gardiner BB, Song S, Harliwong I, Idrisoglu S, Nourse C, Nourbakhsh E, Manning S, Wani S, Gongora M, Pajic M, Scarlett CJ, Gill AJ, Pinho AV, Rooman I, Anderson M, Holmes O, Leonard C, Taylor D, Wood S, Xu Q, Nones K, Fink JL, Christ A, Bruxner T, Cloonan N, Kolle G, Newell F, Pinese M, Mead RS, Humphris JL, Kaplan W, Jones MD, Colvin EK, Nagrial AM, Humphrey ES, Chou A, Chin VT, Chantrill LA, Mawson A, Samra JS, Kench JG, Lovell JA, Daly RJ, Merrett ND, Toon C, Epari K, Nguyen NQ, Barbour A, Zeps N, Australian Pancreatic Cancer Genome I, Kakkar N, Zhao F, Wu YQ, Wang M, Muzny DM, Fisher WE, Brunicki FC, Hodges SE, Reid JG, Drummond J, Chang K, Han Y, Lewis LR, Dinh H, Buhay CJ, Beck T, Timms L, Sam M, Begley K, Brown A, Pai D, Panchal A, Buchner N, De Borja R, Denroche RE, Yung CK, Serra S, Onetto N, Mukhopadhyay D, Tsao MS, Shaw PA, Petersen GM, Gallinger S, Hruban RH, Maitra A, Iacobuzio-Donahue CA, Schulick RD, Wolfgang CL, Morgan RA, Lawlor RT, Capelli P, Corbo V, Scardoni M, Tortora G, Tempero MA, Mann KM, Jenkins NA, Perez-Mancera PA, Adams DJ, Largaespada DA, Wessels LF, Rust AG, Stein LD,

Tuveson DA, Copeland NG, Musgrove EA, Scarpa A, Eshleman JR, Hudson TJ, Sutherland RL, Wheeler DA, Pearson JV, McPherson JD, Gibbs RA and Grimmond SM. (2012) Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. *Nature*; 491: 399-405.

Bidard FC, Huguet F, Louvet C, Mineur L, Bouche O, Chibaudel B, Artru P, Desseigne F, Bachet JB, Mathiot C, Pierga JY and Hammel P. (2013) Circulating tumor cells in locally advanced pancreatic adenocarcinoma: the ancillary CirCe 07 study to the LAP 07 trial. *Ann Oncol*; 24: 2057-2061.

Bloomston M, Frankel WL, Petrocca F, Volinia S, Alder H, Hagan JP, Liu CG, Bhatt D, Taccioli C, Croce CM. (2007) MicroRNA expression patterns to differentiate pancreatic adenocarcinoma from normal pancreas and chronic pancreatitis. *JAMA*; 297: 1901-1908.

Bloomston M, Zhou JX, Rosemurgy AS, Frankel W, Muro-Cacho CA, Yeatman TJ. (2006) Fibrinogen gamma overexpression in pancreatic cancer identified by large-scale proteomic analysis of serum samples. *Cancer Res* 66: 2592 – 2599

Boscher C, Dennis JW, Nabi IR. (2011) Glycosylation, galectins and cellular signaling. *Curr. Opin. Cell Biol.* 23, 383–392.

Bosetti C, Bertuccio P, Malvezzi M, Levi F, Chatenoud L, Negri E, La Vecchia C. (2013) Cancer mortality in Europe, 2005-2009, and an overview of trends since 1980. *Ann Oncol*, 24(10):2657–71. doi: 10.1093/annonc/mdt301.

Bracci PM, Zhou M, Young S and Wiemels J. (2012) Serum autoantibodies to pancreatic cancer antigens as biomarkers of pancreatic cancer in a San Francisco Bay Area case-control study. *Cancer*; 118: 5384-5394.

Brand RE, Nolen BM, Zeh HJ, Allen PJ, Eloubeidi MA, Goldberg M, Elton E, Arnoletti JP, Christein JD, Vickers SM, Langmead CJ, Landsittel DP, Whitcomb DC, Grizzle WE and Lokshin AE. (2011) Serum biomarker panels for the detection of pancreatic cancer. *Clin Cancer Res*; 17: 805-816.

Bray I, Gunnell D, Holly JM, Middleton N, Smith GD, Martin RM. (2006) Associations of childhood and adulthood height and the components of height with insulin-like growth factor levels in adulthood: a 65-year follow-up of the Boyd Orr cohort. *J Clin Endocrinol Metab*; 91: 1382–9. doi: 10.1210/jc.2005-1722

Brennan DD, Zamboni GA, Raptopoulos VD, Kruskal JB. (2007) Comprehensive preoperative assessment of pancreatic adenocarcinoma with 64-section volumetric CT. *Radiographics*; 27: 1653-1666.

Brierley JD, Gospodarowicz MK, Wittekind C, editors. UICC. TNM (2017) Classification of malignant tumors. 8th ed.; Wiley-Blackwell: Hoboken, NJ, USA

Brockhausen I. (2006) Mucin-type O-glycans in human colon and breast cancer: glycodynamics and functions. *EMBO Rep.* 7, 599–604.

Bunger S, Laubert T, Roblick UJ and Habermann JK. (2011) Serum biomarkers for improved diagnostic of pancreatic cancer: a current overview. *J Cancer Res Clin Oncol*; 137: 375-389.

Calle EE, Kaaks R. (2004) Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer*; 4: 579-91.

Cao Z, Liu C, Xu J, You L, Wang C, Lou W, Sun B, Miao Y, Liu X, Wang X, Zhang T, Zhao Y. (2016) Plasma microRNA panels to diagnose pancreatic cancer: results from a multicenter study. *Oncotarget*; 7: 41575-41583.

Capello M, Bantis LE, Scelo G, Zhao Y, Li P, Dhillon DS, Patel NJ, Kundnani DL, Wang H, Abbruzzese JL, Maitra A, Tempero MA, Brand R, Firpo MA, Mulvihill SJ, Katz MH, Brennan P, Feng Z, Taguchi A, Hanash SM. (2017) Sequential Validation of Blood-Based Protein Biomarker Candidates for Early-Stage Pancreatic Cancer. *J Natl Cancer Inst*; 109. PMID: 28376157 DOI: 10.1093/jnci/ djw266

Capello M, Cappello P, Linty FC, Chiarle R, Sperduti I, Novarino A, Salacone P, Mandili G, Naccarati A, Sacerdote C, Beghelli S, Bersani S, Barbi S, Bassi C, Scarpa A, Nistico P, Giovarelli M, Vineis P, Milella M and Novelli F. (2013) Auto-antibodies to Ezrin are an early sign of pancreatic cancer in humans and in genetically engineered mouse models. *J Hematol Oncol*; 6: 67. doi: 10.1186/1756-8722-6-67.

Caponi S, Funel N, Frampton AE, Mosca F, Santarpia L, Van der Velde AG, Jiao LR, De Lio N, Falcone A, Kazemier G, Meijer GA, Verheul HM, Vasile E, Peters GJ, Boggi U and Giovannetti E. (2013) The good, the bad and the ugly: a tale of miR- 101, miR- 21 and miR-155 in pancreatic intra- ductal papillary mucinous neoplasms. *Ann Oncol*; 24: 734-741.

Carvalhoc AS, Harduin-Lepers A, Magalhães A, Machado E, Mendes N, Costa LT, Matthiesen R, Almeida R, Costa J, Reis CA. (2010) Differential expression of  $\alpha$ -2,3-sialyltransferases and  $\alpha$ -1,3/4-fucosyltransferases regulates the levels of sialyl Lewis a and sialyl Lewis x in gastrointestinal carcinoma cells. *Int. J. Biochem. Cell Biol.* 42, 80–89. doi: 10.1016/j.biocel.2009.09.010.

Castells A, Puig P, Mora J, Boadas J, Boix L, Urgell E, Sole M, Capella G, Lluís F, Fernandez- Cruz L, Navarro S and Farre A. (1999) K-ras mutations in DNA extracted

from the plasma of patients with pancreatic carcinoma: diagnostic utility and prognostic significance. *J Clin Oncol*; 17: 578-584.

Chang JC, Kundranda M. (2017) Novel Diagnostic and Predictive Biomarkers in Pancreatic Adenocarcinoma. *Int J Mol Sci*; 18. PMID: 28335509 DOI: 10.3390/ijms18030667

Chang ST, Zahn JM, Horecka J, Kunz PL, Ford JM, Fisher GA, Le QT, Chang DT, Ji H and Koong AC. (2009) Identification of a biomarker panel using a multiplex proximity ligation assay improves accuracy of pancreatic cancer diagnosis. *J Transl Med*; 7: 105.

Charrier A, Chen R, Chen L, Kemper S, Hattori T, Takigawa M, Brigstock DR. (2014) Connective tissue growth factor (CCN2) and microRNA-21 are components of a positive feedback loop in pancreatic stellate cells (PSC) during chronic pancreatitis and are exported in PSC-derived exosomes. *J Cell Commun Signal*; 8: 147-156.

Chen J, Yang R, Lu Y, Xia Y, Zhou H. (2012) Diagnostic accuracy of endoscopic ultrasound-guided fine-needle aspiration for solid pancreatic lesion: a systematic review. *J Cancer Res Clin Oncol*; 138: 1433-1441.

Chen S, LaRoche T, Hamelinck D, Bergsma D, Brenner D, Simeone D, Brand RE, Haab BB. (2007) Multiplexed analysis of glycan variation on native proteins captured by antibody microarrays. *Nat Methods*; 4: 437-444. PMID: 17417647 DOI: 10.1038/nmeth1035

Ching CK, Rhodes JM. (1989) Enzyme-linked PNA lectin binding assay compared with CA19-9 and CEA radioimmunoassay as a diagnostic blood test for pancreatic cancer. *British Journal of Cancer* 59(6):949–953. DOI 10.1038/bjc.1989.202.

Cohen J.D, Javed AA, Thoburn C, Wong F, Tie J, Gibbs P, Schmidt CM, Yip-Schneider MT, Allen PJ, Schattner M, Brand Re, Singhi AD, Peterson GM, Hong SM, Kim SC, Falconi M, Doglioni C, Weiss MJ, Ahuja M, He J, Makary MA, Maitra A, Hanash SM, Dal Molin M, Wang Y, Li L, Ptak J, Dobbryn L, Schaefer J, Silliman N, Popoli M, Goggins MG, Hruban RH, Wolfgang CL, Klein AP, Tomasetti C, Papadopoulos N, Kinzler KW, Voglestein B, Lennon AM. (2017) Combined circulating tumor DNA and protein biomarker-based liquid biopsy for the earlier detection of pancreatic cancers. *Proc. Natl. Acad. Sci. USA*, 114, 10202–10207.doi: 10.1073/pnas.1704961114.

Costello E, Greenhalf W and Neoptolemos JP. (2012) New biomarkers and targets in pancreatic cancer and their application to treatment. *Nat Rev Gastroenterol Hepatol*; 9: 435-444.



Crawford HC, Scoggins CR, Washington MK, Matrisian LM, Leach SD. (2002) Matrix metalloproteinase-7 is expressed by pancreatic cancer precursors and regulates acinar-to-ductal metaplasia in exocrine pancreas. *J Clin Invest* 109: 1437 – 1444.

Croci DO, Cerliani JP, Dalotto-Moreno T, Méndez-Huergo SP, Mascanfroni ID, Dergan-Dylon S, Toscano MA, Caramelo JJ, García-Vallejo JJ, Ouyang J, Mesri EA, Junttila MR, Bais C, Shipp MA, Salatino M, Rabinovich GA. (2014) Glycosylation-dependent lectin- receptor interactions preserve angiogenesis in anti- VEGF refractory tumors. *Cell* 156, 744–758. doi: 10.1016/j.cell.2014.01.043.

Cui Q, Lewis IA, Hegeman AD, Anderson ME, Li J, Schulte CF, Westler WM, Eghbalnia HR, Sussman MR and Markley JL. (2008) Metabolite identification via the Madison Metabolomics Consortium Database. *Nat Biotechnol*; 26: 162-164.

Cummings RD. (2009) The repertoire of glycan determinants in the human glycome. *Mol. Biosyst.* 5, 1087–1104.

de-Freitas-Junior JCM, Carvalho S, Dias AM, Oliveira P, Cabral J, Seruca R, Oliveira C, Morgado-Díaz JA, Reis CA, Pinho SS. (2013) Insulin/ IGF-I Signaling Pathways Enhances Tumor Cell Invasion through Bisecting GlcNAc N-glycans Modulation. An Interplay with E-Cadherin. *PLOS ONE* 8(11): e81579. <https://doi.org/10.1371/journal.pone.0081579>

Denkert C, Bucher E, Hilvo M, Salek R, Oresic M, Griffin J, Brockmoller S, Klauschen F, Loibl S, Barupal DK, Budczies J, Iljin K, Nekljudova V and Fiehn O. (2012) Metabolomics of human breast cancer: new approaches for tumor typing and biomarker discovery. *Genome Med*; 4: 37.

Dennis J, Laferte S, Waghorne C, Breitman M, Kerbel R. (1987) Beta 1–6 branching of Asn-linked oligosaccharides is directly associated with metastasis. *Science* 236, 582–585.

Desmetz C, Mange A, Maudelonde T, Solassol J. (2011) Autoantibody signatures: progress and perspectives for early cancer detection. *J Cell Mol Med*; 15: 2013-2024.

Di Gangi IM, Mazza T, Fontana A, Copetti M, Fusilli C, Ippolito A, Mattivi F, Latiano A, Andriulli A, Vrhovsek U and Paziienza V. (2016) Metabolomic profile in pancreatic cancer patients: a consensus-based approach to identify highly discriminating metabolites. *Oncotarget*; 7: 5815- 5829. doi: 10.18632/oncotarget.6808.

Di Lella S, Sunblad V, Cerliani JP, Guardia CM, Estrin DA, Vasta GR, Rabinovich GA. (2011) When galectins recognize glycans: from biochemistry to physiology and back again. *Biochemistry* 50, 7842–7857. doi: 10.1021/bi201121m

Diamandis EP. (2010) Cancer biomarkers: can we turn recent failures into success? *J Natl Cancer Inst*; 102: 1462-1467. PMID: 20705936 DOI: 10.1093/jnci/djq306.

Diaz LA Jr, Bardelli A. (2014) Liquid biopsies: genotyping circulating tumor DNA. *J Clin Oncol*; 32: 579-586.

Dibble EH, Karantanis D, Mercier G, Peller PJ, Kachnic LA, Subramaniam RM. (2012) PET/CT of cancer patients: part 1, pancreatic neoplasms. *AJR Am J Roentgenol*; 199: 952-967. PMID: 23096166 DOI: 10.2214/AJR.11.8182

Ding G, Zhou L, Qian Y, Fu M, Chen J, Chen J, Xiang J, Wu Z, Jiang G, Cao L. (2015) Pancreatic cancer-derived exosomes transfer miRNAs to dendritic cells and inhibit RFXAP expression via miR-212-3p. *Oncotarget*; 6:29877-88. doi: 10.18632/oncotarget.4924.

Djebali S, Davis CA, Merkel A, Dobin A, Lassmann T, Mortazavi A, Tanzer A, Lagarde J, Lin W, Schlesinger F, Xue C, Marinov GK, Khatun J, Williams BA, Zaleski C, Rozowsky J, Roder M, Kokocinski F, Abdelhamid RF, Alioto T, Antoshechkin I, Baer MT, Bar NS, Batut P, Bell K, Bell I, Chakraborty S, Chen X, Chrast J, Curado J, Derrien T, Drenkow J, Dumais E, Dumais J, Duttagupta R, Falconnet E, Fastuca M, Fejes-Toth K, Ferreira P, Foissac S, Fullwood MJ, Gao H, Gonzalez D, Gordon A, Gunawardena H, Howald C, Jha S, Johnson R, Kapranov P, King B, Kingswood C, Luo OJ, Park E, Persaud K, Preall JB, Ribeca P, Risk B, Robyr D, Sammeth M, Schaffer L, See LH, Shahab A, Skancke J, Suzuki AM, Takahashi H, Tilgner H, Trout D, Walters N, Wang H, Wrobel J, Yu Y, Ruan X, Hayashizaki Y, Harrow J, Gerstein M, Hubbard T, Reymond A, Antonarakis SE, Hannon G, Giddings MC, Ruan Y, Wold B, Carninci P, Guigo R and Gingeras TR. (2012) Landscape of transcription in human cells. *Nature*; 489:101-108.

Drabik A, Bodzon-Kulakowska A, Suder P, Silberring J, Kulig J, Sierzega M. (2017) Glycosylation Changes in Serum Proteins Identify Patients with Pancreatic Cancer. *J Proteome Res*; 16: 1436-1444. PMID: 28244758

DOI: 10.1021/acs.jproteome.6b00775

du Rieu MC, Torrisani J, Selves J, Al Saati T, Souque A, Dufresne M, Tsongalis GJ, Suriawinata AA, Carrere N, Buscail L and Cordelier P. (2010) MicroRNA-21 is induced early in pancreatic ductal adenocarcinoma precursor lesions. *Clin Chem*; 56: 603-612.

Duan W, Chen K, Jiang Z, Chen X, Sun L, Li J, Lei J, Xu Q, Ma J, Li X, Han L, Wang Z, Wu Z, Wang F, Wu E, Ma Q, Ma Z. (2017) Desmoplasia suppression by metformin-

mediated AMPK activation inhibits pancreatic cancer progression. *Cancer Lett.*;385:225–33. doi: 10.1016/j.canlet.2016.10.019.

Duell EJ. (2012) Epidemiology and potential mechanisms of tobacco smoking and heavy alcohol consumption in pancreatic cancer. *Molecular Carcinogenesis*; 51: 40-52.

Erkan M, Michalski CW, Rieder S, Reiser-Erkan C, Abiatari I, Kolb A, Giese NA, Esposito I, Friess H, Kleeff J. (2008) The activated stroma index is a novel and independent prognostic marker in pancreatic ductal adenocarcinoma. *Clin Gastroenterol Hepatol*; 6(10):1155–61.

Erkan M. (2013) Understanding the stroma of pancreatic cancer: co-evolution of the microenvironment with epithelial carcinogenesis. *J Pathol* 231: 4 – 7.

Faca VM, Song KS, Wang H, Zhang Q, Krasnoselsky AL, Newcomb LF, Plentz RR, Gurumurthy S, Redston MS, Pitteri SJ, Pereira-Faca SR, Ireton RC, Katayama H, Glykhova V, Phanstiel D, Brenner DE, Anderson MA, Misek D, Scholler N, Urban ND; Barnett MJ, Edelstein C, Goodman GE, Thornquist MD, McIntosh MW, DePinho RA, Bardeesy N, Hanash SM. (2008) A mouse to human search for plasma proteome changes associated with pancreatic tumor development. *PLoS Med* 5: e123. doi: 10.1371/journal.pmed.0050123.

Feng Z, Kagan J, Pepe M, Wagner P, Thornquist M, Rinaudo JA; Dahlfren J, Krueger K, Zheng Y, Patriotis C, Huang Y, Thompson I, Srivastava S. (2013) The Early Detection Research Network's Specimen reference sets: paving the way for rapid evaluation of potential biomarkers. *Clin Chem.*; 59:68–74.

doi: [10.1373/clinchem.2012.185140](https://doi.org/10.1373/clinchem.2012.185140)

Ferlay J, Shin HR, Bray F, (2010) 'GLOBOCAN 2008, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 10 [Internet]. Lyon, France: International Agency for Research on Cancer; Available from: <http://globocan.iarc.fr> 2010

Ferlay J, Soerjomataram I, Ervik M, GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11. Lyon, France: International Agency for Research on Cancer; 2013. Available from: <http://globocan.iarc.fr>, accessed January 2019.

Ferrer-Batallé M, Llop E, Ramírez M, Aleixandre RN, Saez M, Comet J, de Llorens R, Peracaula R. (2017) Comparative Study of Blood- Based Biomarkers,  $\alpha$ 2,3-Sialic Acid PSA and PHI, for High-Risk Prostate Cancer Detection. *Int J Mol Sci*; 18. PMID: 28420168 DOI: 10.3390/ijms18040845

Ferri MJ, Saez M, Figueras J, Fort E, Sabat M, López-Bem S, de Llorens R, Aleixandre RN, Peracaula R. (2016) Improved Pancreatic Adenocarcinoma Diagnosis in Jaundiced and Non- Jaundiced Pancreatic Adenocarcinoma Patients through the Combination of Routine Clinical Markers Associated to Pancreatic Adenocarcinoma Pathophysiology. PLoS One; 11: e0147214. doi: 10.1371/journal.pone.0147214.

Fiedler GM, Leichtle AB, Kase J, Baumann S, Ceglarek U, Felix K, Conrad T, Witzigmann H, Weimann A, Schutte C, Hauss J, Büchler M, Thiery J. (2009) Serum peptidome profiling revealed platelet factor 4 as a potential discriminating Peptide associated with pancreatic cancer. Clin Cancer Res 15: 3812 – 3819. doi: 10.1158/1078-0432.CCR-08-2701.

Fleischhacker M, Schmidt B. (2007) Circulating nucleic acids (CNAs) and cancer - a survey. Biochim Biophys Acta; 1775: 181-232.

Flippot R, Malouf GG, Su X, Mouawad R, Spano JP, Khayat D. (2016) Cancer subtypes classification using long non-coding RNA. Oncotarget; 7: 54082-54093.

Freeze HH. (2013) Understanding human glycosylation disorders: biochemistry leads the charge. J. Biol. Chem. 288, 6936–6945.

Friedenreich CM, Neilson HK, Lynch BM. (2010) State of the epidemiological evidence on physical activity and cancer prevention. Eur J Cancer; 46: 2593–604.

Fu Z, Chen C, Zhou Q, Wang Y, Zhao Y, Zhao X, Li W, Zheng S, Ye H, Wang L, He Z, Lin Q, Li Z, Chen R. (2017) LncRNA HOTTIP modulates cancer stem cell properties in human pancreatic cancer by regulating HOXA9. Cancer Lett; 410: 68-81.

Fujita N, Noda Y, Kobayashi G, Go MD, Kimura K, Katsumi MD, Ito K, Kei MD. (2004) Endoscopic approach to early diagnosis of pancreatic cancer. Pancreas;28:279-81.

Fukushima N, Walter KM, Uek T, Sato N, Matsubayashi H, Cameron JL, Hruban RH, Canto M, Yeo CJ, Goggins M. (2003) Diagnosing pancreatic cancer using methylation specific PCR analysis of pancreatic juice. Cancer Biol Ther;2:78-83. 35.

Fuster, M, Esko, J. (2005) The sweet and sour of cancer: glycans as novel therapeutic targets. Nat. Rev. Cancer 5, 526–542.

Füzéry AK, Levin J, Chan MM, Chan DW. (2013) Translation of proteomic biomarkers into FDA approved cancer diagnostics: issues and challenges. Clin Proteomics; 10: 13. PMID: 24088261 DOI: 10.1186/1559-0275-10-13

Gao Y, Zhu Y, Zhang Z, Zhang C, Huang X, Yuan Z. (2016) Clinical significance of pancreatic circulating tumor cells using combined negative enrichment and immunostaining-fluorescence in situ hybridization. *J Exp Clin Cancer Res*; 35: 66.

Ghaneh P, Costello E, Neoptolemos JP. (2007) Biology and management of pancreatic cancer. *Gut*; 56: 1134-1152.

Gill DJ, Chia J, Senewiratne J, Bard F. (2010) Regulation of O-glycosylation through Golgi-to-ER relocation of initiation enzymes. *J. Cell Biol.* 189, 843–858.

Giménez E, Balmaña M, Figueras J, Fort E, de Bolós C, Sanz- Nebot V, Peracaula R, Rizzi A. (2015) Quantitative analysis of N-glycans from human alfa-acid-glycoprotein using stable isotope labeling and zwitterionic hydrophilic interaction capillary liquid chromatography electrospray mass spectrometry as tool for pancreatic disease diagnosis. *Anal Chim Acta*; 866: 59-68. PMID: 25732693 DOI: 10.1016/j.aca.2015.02.008

Giulietti M, Righetti A, Principato G, Piva F. (2018) LncRNA co-expression network analysis reveals novel biomarkers for pancreatic cancer, *Carcinogenesis*, Volume 39, Issue 8, Pages 1016–1025. doi: 10.1093/carcin/bgy069

Goh SK, Gold G, Christophi C, Muralidharan V. (2017) Serum carbohydrate antigen 19-9 in pancreatic adenocarcinoma: a mini review for surgeons. *ANZ J Surg*; 87: 987-992. PMID: 28803454 DOI: 10.1111/ans.14131

Gold DV, Gaedcke J, Ghadimi BM, Goggins M, Hruban RH, Liu M, Newsome G, Golden-berg DM. (2013) PAM4 enzyme immunoassay alone and in combination with CA 19-9 for the detection of pancreatic adenocarcinoma. *Cancer*; 119: 522-528.

Gold DV, Modrak DE, Ying Z, Cardillo TM, Sharkey RM, Goldenberg DM. (2006) New MUC1 serum immunoassay differentiates pancreatic cancer from pancreatitis. *J Clin Oncol*; 24: 252-258.

Gomes C, Osório H, Pinto MT, ampos D, Oliveira MJ, Reis CA. (2013) Expression of ST3GAL4 leads to SLe<sup>x</sup> expression and induces c-Met activation and an invasive phenotype in gastric carcinoma cells. *PLOS ONE* 8(6): e66737. <https://doi.org/10.1371/journal.pone.0066737>

Greer JB, Whitcomb DC, Brand RE. (2007) Genetic predisposition to pancreatic cancer: a brief review. *Am J Gastroenterol*; 102: 2564-2569.

Gu J, Sato Y, Kariya Y, Isaji T, Taniguchi N, Fukuda T. (2009) A mutual regulation between cell-cell adhesion and N-glycosylation: implication of the bisecting GlcNAc for biological functions. *J. Proteome Res.* 8, 431–435. doi: 10.1021/pr800674g.

Gunnell D, Okasha M, Smith GD, Oliver SE, Sandhu J, Holly JM. (2001) Height, leg length, and cancer risk: a systematic review. *Epidemiol Rev*; 23: 313–42.

Haab BB, Huang Y, Balasenthil S, Partyka K, Tang H, Anderson M, Allen P, Sasson A, Zeh H, Kaul K, Kletter D, Ge S, Bern M, Kwon R, Blasutig I, Srivastava S, Frazier ML, Sen S, Hollingsworth MA, Rinaudo JA, Killary AM, Brand RE. (2015) Definitive Characterization of CA 19-9 in Resectable Pancreatic Cancer Using a Reference Set of Serum and Plasma Specimens. *PloS One.*; 10: e0139049. <https://doi.org/10.1371/journal.pone.0139049>

Haab BB, Porter A, Yue T, Li L, Scheiman J, Anderson MA, Barnes D, Schmidt CM, Feng Z, Simeone DM. (2010) Glycosylation variants of mucins and CEACAMs as candidate biomarkers for the diagnosis of pancreatic cystic neoplasms. *Ann Surg*; 251: 937-945. PMID: 20395854 DOI: 10.1097/SLA.0b013e3181d7738d]

Hage DS, Anguizola JA, Bi C, Li R, Matsuda R, Papastavros E, Pfaunmiller E, Vargas J, Zheng X. (2012) Pharmaceutical and biomedical applications of affinity chromatography: recent trends and developments. *Journal of Pharmaceutical and Biomedical Analysis* 69:93–105. DOI 10.1016/j.jpba.2012.01.004.

Hakama M, Louhivuori K. (1988) A screening program for cervical cancer that worked. *Cancer Surv.*; 7:403–416.

Hakomori S, Kannagi R. (1983) Glycosphingolipids as tumor-associated and differentiation markers. *J. Natl Cancer Inst.* 71, 231–251.

Hakomori S. (2002) Glycosylation defining cancer malignancy: new wine in an old bottle. *Proc. Natl Acad. Sci. USA* 99, 10231–10233.

Hakomori SI, Murakami WT. (1968) Glycolipids of hamster fibroblasts and derived malignant- transformed cell lines. *Proc. Natl Acad. Sci. USA* 59, 254–261.

Hanoun N, Delpu Y, Suriawinata AA, Bournet B, Bureau C, Selves J, Tsongalis GJ, Dufresne M, Buscail L, Cordelier P, Torrisani J. (2010) The silencing of microRNA 148a production by DNA hypermethylation is an early event in pancreatic carcinogenesis. *Clin Chem*;56:1107-18. DOI: 10.1373/clinchem.2010.144709

Hart GW. (2016) Myriad Roles of Glycans in Biology. *J Mol Biol*; 428: 3147-3149. PMID: 27342936 DOI: 10.1016/j.jmb.2016.06.010

Hashim O, Jayapalan J, Lee C. (2017), Lectins: an effective tool for screening of potential cancer biomarkers. *PeerJ* 5:e3784. DOI 10.7717/peerj.3784

Hatano K, Miyamoto Y, Nonomura N, Kaneda Y. (2011) Expression of gangliosides, GD1a, and sialyl paragloboside is regulated by NF- $\kappa$ B- dependent transcriptional control of  $\alpha$ 2,3-sialyltransferase I, II, and VI in human castration-resistant prostate cancer cells. *Int.J. Cancer* 129, 1838–1847.

Helenius A, Aebi M. (2001) Intracellular functions of N-linked glycans. *Science* 291, 2364–2369.

Hengstler JG, Foth H, Gebel T, Kramer PJ, Lilienblum W, Schweinfurth H, Völkel W, Wollin KM, Gundert-Remy U. (2011) Critical evaluation of key evidence on the human health hazards of exposure to bisphenol A. *Crit Rev Toxicol*, 41:263–291. doi: 10.3109/10408444.2011.558487.

Hernandez YG, Lucas AL. (2016) MicroRNA in pancreatic ductal adenocarcinoma and its precursor lesions. *World J Gastrointest Oncol*;8:18-29.

Herreros-Villanueva M, Bujanda L. (2016) Non-invasive biomarkers in pancreatic cancer diagnosis: what we need versus what we have, *Ann Transl Med.*;4(7):134.

Hewitt MJ, McPhail MJ, Possamai L, Dhar A, Vlavianos P, Monahan KJ. (2012) EUS-guided FNA for diagnosis of solid pancreatic neoplasms: a meta-analysis. *Gastrointest Endosc*; 75: 319-331.

Hidalgo M. (2010) Pancreatic cancer. *N Engl J Med*; 362: 1605-1617.

Hirabayashi J, Kuno A, Tateno H. (2011) Lectin-based structural glycomics: a practical approach to complex glycans. *Electrophoresis* 32(10):1118–1128. DOI 10.1002/elps.201000650.

Hong SH, Misek DE, Wang H, Puravs E, Giordano TJ, Greenson JK, Brenner DE, Simeone DM, Logsdon CD, Hanash SM. (2004) An autoantibody-mediated immune response to calreticulin isoforms in pancreatic cancer. *Cancer Res*; 64: 5504-5510. DOI: 10.1158/0008-5472.CAN-04-0077

Hong SH, Misek DE, Wang H, Puravs E, Hinderer R, Giordano TJ, Greenson JK, Brenner DE, Simeone DM, Logsdon CD, Hanash SM. (2006) Identification of a specific vimentin isoform that induces an antibody response in pancreatic cancer. *Biomark Insights*; 1: 175-183.

Hoshino A, Costa-Silva B, Shen TL, Rodrigues G, Hashimoto A, Tesic Mark M, Molina H, Koh- saka S, Di Giannatale A, Ceder S, Singh S, Williams C, Soplop N, Uryu K,

Pharmer L, King T, Bojmar L, Davies AE, Ararso Y, Zhang T, Zhang H, Hernandez J, Weiss JM, Dumont-Cole VD, Kramer K, Wexler LH, Narendran A, Schwartz GK, Healey JH, Sandstrom P, Labori KJ, Kure EH, Grandgenett PM, Hollingsworth MA, de Sousa M, Kaur S, Jain M, Mallya K, Batra SK, Jarnagin WR, Brady MS, Fodstad O, Muller V, Pantel K, Minn AJ, Bissell MJ, Garcia BA, Kang Y, Rajasekhar VK, Ghajar CM, Matei I, Peinado H, Bromberg J, Lyden D. (2015) Tumor exosome integrins determine organotropic metastasis. *Nature*; 527: 329-335.

Hruban RH, Pitman MB, Klimstra DS. (2007). *Tumors of the Pancreas (AFIP Atlas of Tumor Pathology Series 4, Fascicle 6)*. Washington, DC: American Registry of Pathology and Armed Forces Institute of Pathology.

Hu S, Wong DT. (2009) Lectin microarray. *Proteomics Clinical Applications* 3(2):148–154. DOI 10.1002/prca.200800153.

Huang Z, Liu F. (2014) Diagnostic value of serum carbohydrate antigen 19-9 in pancreatic cancer: a meta-analysis. *Tumour Biol*; 35: 7459-7465.

Hur C, Tramontano A, Dowling E, Brooks G, Jeon, A, Brugge W, Gazelle G, Kong C, Pandharipande P. (2016) Early Pancreatic Ductal Adenocarcinoma Survival is Dependent on Size: Positive Implications for Future Targeted Screening, *MPH Pancreas*; 45(7): 1062–1066.

Hursting SD, Lavigne JA, Berrigan D, Perkins SN, Barret JC. (2003) 'Calorie restriction, aging, and cancer prevention: mechanisms of action and applicability to humans. *Annu Rev Med*; 54: 131-52.

Husemann Y, Geigl JB, Schubert F, Musiani P, Meyer M, Burghart E, Forni G, Eils R, Fehm T, Riethmuller G, Klein CA. (2008) Systemic spread is an early step in breast cancer. *Cancer Cell*; 13: 58-68.

Hwang RF, Moore T, Arumugam T, Ramachandran V, Amos KD, Rivera A, Ji B, Evans DB, Logsdon CD. (2008) Cancer-associated stromal fibroblasts promote pancreatic tumor progression. *Cancer Res*; 68(3):918–26.

IARC (1988). Alcohol drinking. *IARC Monogr Eval Carcinog Risks Hum*, 44:1– 378. PMID:3236394

Imamura T, Komatsu S, Ichikawa D, Kawaguchi T, Miyamae M, Okajima W, Ohashi T, Arita T, Konishi H, Shiozaki A, Morimura R, Ikoma H, Okamoto K, Otsuji E. (2016) Liquid biopsy in patients with pancreatic cancer: Circulating tumor cells and cell-free nucleic acids. *World J Gastroenterol*; 22: 5627-5641. PMID: 27433079 DOI: 10.3748/wjg.v22.i25.5627



International Agency for Research on Cancer, World cancer report (2014)

Iorio MV, Croce CM. (2009) MicroRNAs in cancer: small molecules with a huge impact. *J Clin Oncol*;27:5848-56.

ISD Scotland, Detect Cancer Early Staging Data. Scotland: ISD; 2016.

Iwanicki-Caron I, Basile P, Toure E, Antonietti M, Lecleire S, Di Fiore A, Oden-Gangloff A, Blanchard F, Lemoine F, Di Fiore F, Sabourin JC and Michel P. (2013) Usefulness of circulating tumor cell detection in pancreatic adenocarcinoma diagnosis. *Am J Gastroenterol*; 108: 152- 155.

Jiang Z, Zhou C, Cheng L, Yan B, Chen K, Chen X, Zong L, Lei J, Duan W, Xu Q, Li X, Wang Z, Ma Q, Ma J. (2018) Inhibiting YAP expression suppresses pancreatic cancer progression by disrupting tumor-stromal interactions, *Journal of Experimental & Clinical Cancer Research*, 37:69

Jiao F, Hu H, Yuan C, Wang L, Jiang W, Jin Z, Guo Z, Wang L. (2014) Elevated expression level of long noncoding RNA MALAT-1 facilitates cell growth, migration and invasion in pancreatic cancer. *Oncol Rep*; 32: 2485-2492.

Jiao L, Zhu J, Hassan MM, Evans DB, Abbruzzese JL, Li D. (2007) K-ras mutation and p16 and preproenkephalin promoter hypermethylation in plasma DNA of pancreatic cancer patients: in relation to cigarette smoking. *Pancreas*; 34: 55-62.

Jones S, Zhang X, Parsons DW, Lin JC, Leary RJ, Angenendt P, Mankoo P, Carter H, Kamiyama H, Jimeno A, Hong SM, Fu B, Lin MT, Calhoun ES, Kamiyama M, Walter K, Nikolskaya T, Nikolsky Y, Hartigan J, Smith DR, Hidalgo M, Leach SD, Klein AP, Jaffee EM, Goggins M, Maitra A, Iacobuzio-Donahue C, Eshleman JR, Kern SE, Hruban RH, Karchin R, Papadopoulos N, Parmigiani G, Vogelstein B, Velculescu VE, Kinzler KW. (2008) Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. *Science*; 321: 1801- 1806.

Kahlert C, Melo SA, Protopopov A, Tang J, Seth S, Koch M, Zhang J, Weitz J, Chin L, Futreal A, Kalluri R. (2014) Identification of double-stranded genomic DNA spanning all chromosomes with mutated KRAS and p53 DNA in the serum exosomes of patients with pancreatic cancer. *J Biol Chem*;289:3869-75. doi: 10.1074/jbc.C113.532267.

Kailemia MJ, Park D, Lebrilla CB. (2017) Glycans and glycoproteins as specific biomarkers for cancer. *Anal Bioanal Chem*; 409: 395-410. PMID: 27590322 DOI: 10.1007/s00216-016-9880-6

Kailemia MJ, Xu G, Wong M, Li Q, Goonatilleke E, Leon F, Lebrilla CB. (2018) Recent Advances in the Mass Spectrometry Methods for Glycomics and Cancer. *Anal Chem*; 90: 208-224. PMID: 29049885 DOI: 10.1021/acs.analchem.7b04202

Kakugawa, Y, Wada T, Yamagushi K, Yamanami H, Ouchi K, Sato I, Miyagi T. (2002) Up-regulation of plasma membrane- associated ganglioside sialidase (Neu3) in human colon cancer and its involvement in apoptosis suppression. *Proc. Natl Acad. Sci. USA* 99, 10718–10723. <https://doi.org/10.1073/pnas.152597199>

Kanda M, Matthaei H, Wu J, Hong SM, Yu J, Borges M, Hruban RH, Maitra A, Kinzler K, Vogelstein B, Goggins M. (2012) Presence of somatic mutations in most early-stage pancreatic intraepithelial neoplasia. *Gastroenterology*; 142: 730-733, e739.

Kannagi R, Yin J, Miyazaki K, Izawa M. (2008) Current relevance of incomplete synthesis and neo-synthesis for cancer-associated alteration of carbohydrate determinants—Hakomori's concepts revisited. *Biochim. Biophys. Acta* 1780, 525–531.

Kaur S, Baine MJ, Jain M, Sasson AR, Batra SK. (2012) Early diagnosis of pancreatic cancer: challenges and new developments. *Biomark Med*; 6: 597-612.

Kaur S, Smith LM, Patel A, Menning M, Watley DC, Malik SS, Krishn SR, Mallya K, Aithal A, Sasson AR, Johansson SL, Jain M, Singh S, Guha S, Are C, Raimondo M, Hollingsworth MA, Brand RE, Batra SK. (2017) A Combination of MUC5AC and CA19-9 Improves the Diagnosis of Pancreatic Cancer: A Multicenter Study. *Am J Gastroenterol*. 2017 Jan;112(1):172-183. doi: 10.1038/ajg.2016.482. Epub 2016 Nov 15.

Kellokumpu S, Sormunen R, Kellokumpu I. (2002) Abnormal glycosylation and altered Golgi structure in colorectal cancer: dependence on intra-Golgi pH. *FEBS Lett*. 516, 217–224.

Keri RA, Ho SM, Hunt PA, Knudsen KE, Soto AM, Prins GS. (2007) An evaluation for the carcinogenic activity of bisphenol A. *Reprod Toxicol*, 24:240–252. <http://dx.doi.org/10.1016/j.reprotox.2007.06.008> PMID:17706921

Kim J, Bamlet WR, Oberg AL, Chaffee KG, Donahue G, Cao XJ, Chari S, Garcia BA, Petersen GM, Zaret KS. (2017) Detection of early pancreatic ductal adenocarcinoma with thrombospondin-2 and CA19-9 blood markers. *Sci. Transl. Med.*, 9, eaah5583. doi: 10.1126/scitranslmed.aah5583.

Kim K, Jutooru I, Chadalapaka G, Johnson G, Frank J, Burghardt R, Kim S, Safe S. (2013) HOTAIR is a negative prognostic factor and exhibits pro-oncogenic activity in pancreatic cancer. *Oncogene*; 32: 1616-1625.

Kinugasa H, Nouse K, Miyahara K, Morimoto Y, Dohi C, Tsutsumi K, Kato H, Matsubara T, Okada H, Yamamoto K. (2015) Detection of K-ras gene mutation by liquid biopsy in patients with pancreatic cancer. *Cancer*; 121: 2271-2280.

Kinupe AB, Ung Y, Ko YJ, Berry SR. (2017) FDG PET/CT in pancreatic cancer staging and management: A retrospective study. *Journal of Clin Oncol*; 35: 464-464. DOI: 10.1200/JCO.2017.35.4\_suppl.464

Kisiel JB, Raimondo M, Taylor WR, Yab TC, Mahoney DW, Sun Z, Middha S, Baheti S, Zou H, Smyrk TC, Boardman LA, Peterson GM, Ahlquist DA. (2015) New DNA Methylation Markers for Pancreatic Cancer: Discovery, Tissue Validation, and Pilot Testing in Pancreatic Juice. *Clin Cancer Res*;21:4473-81. doi: 10.1158/1078-0432.CCR-14-2469

Kitamura N, Guo S, Sato T, Hiraizumi S, Taka J, Ikekita M, Sawada S, Fujisawa H, Furukawa K. (2003) Prognostic significance of reduced expression of beta-N -acetylgalactosaminylated N-linked oligosaccharides in human breast cancer. *International Journal of Cancer* 105:533–541. DOI 10.1002/ijc.11115.

Kleeff J, Korc M, Apte M, La Vecchia C, Johnson CD, Biankin AV, Neale RE, Tempero M, Tuveson DA, Hruban RH, Neoptolemos JP. (2016) Pancreatic cancer. *Nat. Rev. Dis. Prim.*, 2, 16022. doi: 10.1038/nrdp.2016.22.

Klein AP, Wolpin BM, Risch HA, Stolzenberg-Solomon RZ, Mocci E, Zhang M, Canzian F, Childs EJ, Hoskins JW, Jermusyk A, Zhong J, Chen F, Albanes D, Andreotti G, Arslan AA, Babic A, Bamlet WR, Beane-Freeman L, Berndt SI, Blackford A, Borges M, Borgida A, Bracci PM, Brais L, Brennan P, Brenner H, Bueno-de-Mesquita B, Buring J, Campa D, Capurso G, Cavestro GM, Chaffee KG, Chung CC, Cleary S, Cotterchio M, Dijk F, Duell EJ, Foretova L, Fuchs C, Funel N, Gallinger SM, Gaziano JM, Gazouli M, Giles GG, Giovannucci E, Goggins M, Goodman GE, Goodman PJ, Hackert T, Haiman C, Hartge P, Hasan M, Hegyi P, Helzlsouer KJ, Herman J, Holcatova I, Holly EA, Hoover R, Hung RJ, Jacobs EJ, Jamroziak K, Janout V, Kaaks R, Khaw KT, Klein EA, Kogevinas M, Kooperberg C, Kulke MH, Kupcinskis J, Kurtz RJ, Laheru D, Landi S, Lawlor RT, Lee IM, LeMarchand L, Lu L, Malats N, Mambrini A, Mannisto S, Milne RL, Mohelníková-Duchoňová B, Neale RE, Neoptolemos JP, Oberg AL, Olson SH, Orlov I, Pasquali C, Patel AV, Peters U, Pezzilli R, Porta M, Real FX, Rothman N, Scelo G, Sesso HD, Severi G, Shu XO, Silverman D, Smith JP, Soucek P, Sund M, Talar-Wojnarowska R, Tavano F, Thornquist MD, Tobias GS, Van Den Eeden SK, Vashist Y, Visvanathan K, Vodicka P, Wactawski-Wende J, Wang Z, Wentzensen N, White E, Yu H, Yu K, Zeleniuch-Jacquotte A, Zheng W, Kraft P, Li D, Chanock S,

Obazee O, Petersen GM, Amundadottir LT. (2018) Genome-wide meta-analysis identifies five new susceptibility loci for pancreatic cancer. *Nature communications*, 9(1), 556.

Kobayashi T, Nishiumi S, Ikeda A, Yoshie T, Sakai A, Matsubara A, Izumi Y, Tsumura H, Tsuda M, Nishisaki H, Hayashi N, Kawano S, Fujiwara Y, Minami H, Takenawa T, Azuma T and Yoshida M. (2013) A novel serum metabolomics-based diagnostic approach to pancreatic cancer. *Cancer Epidemiol Biomarkers Prev*; 22: 571-579.

Kobold S, Lutkens T, Cao Y, Bokemeyer C, Atanackovic D. (2010) Autoantibodies against tumor- related antigens: incidence and biologic significance. *Hum Immunol*; 71: 643-651.

Kohler BA, Sherman RL, Howlader N, Jemal A, Ryerson AB, Henry KA, Boscoe FP, Cronin KA, Lake A, Noone AM, Henley SJ, Ehemann CR, Anderson RN, Penberthy L. (2015). Annual Report to the Nation on the Status of Cancer, 1975-2011, featuring incidence of breast cancer subtypes by race/ethnicity, poverty, and state. *J Natl Cancer Inst*, 107(6):djv048. doi: 10.1093/jnci/djv048.

Kojima K, Asmellash S, Klug CA, Grizzle WE, Mobley JA, Christein JD. (2008) Applying proteomic-based biomarker tools for the accurate diagnosis of pancreatic cancer. *J Gastrointest Surg* 12: 1683 – 1690.

Komatsu S, Ichikawa D, Miyamae M, Kawaguchi T, Morimura R, Hirajima S, Okajima W, Ohashi T, Imamura T, Konishi H, Shiozaki A, Ikoma H, Okamoto K, Taniguchi H, Otsuji E. (2015) Malignant potential in pancreatic neoplasm; new insights provided by circulating miR-223 in plasma. *Expert Opin Biol Ther*;15:773-85. doi: 10.1517/14712598.2015.1029914.

Kondo S, Sasaki M, Hosoi H, Sakamoto Y, Morizane C, Ueno H, Okusaka, T. (2018) Incidence and risk factors for venous thromboembolism in patients with pretreated advanced pancreatic carcinoma. *Oncotarget*, 9(24), 16883-16890.

Kontro H, Joenväärä S, Haglund C, Renkonen R. (2014) Comparison of sialylated N-glycopeptide levels in serum of pancreatic cancer patients, acute pancreatitis patients, and healthy controls. *Proteomics*; 14: 1713-1723. PMID: 24841998 DOI: 10.1002/pmic.201300270

Koopmann J, Buckhaults P, Brown DA, Zahurak ML, Sato N, Fukushima N, Sokoll LJ, Chan DW, Yeo CJ, Hruban RH, Breit SN, Kinzler KW, Vogelstein B, Goggins M. (2004) Serum macrophage inhibitory cytokine 1 as a marker of pancreatic and other periampullary cancers. *Clin Cancer Res*; 10: 2386-2392.

- Kumamoto K, Goto Y, Sekikawa K, Takenoshita S, Ishida N, Kawakita M, Kannagi R. (2001) Increased expression of UDP- galactose transporter messenger RNA in human colon cancer tissues and its implication in synthesis of Thomsen-Friedenreich antigen and sialyl Lewis A/X determinants. *Cancer Res.* 61, 4620–4627.
- Kuzmanov U, Kosanam H, Diamandis EP. (2013) The sweet and sour of serological glycoprotein tumor biomarker quantification. *BMC Medicine* 11:31. DOI 10.1186/1741-7015-11-31.
- La Thangue N, Kerr D. (2011) Predictive biomarkers: a paradigm shift towards personalized cancer medicine, *Nature Reviews Clinical Oncology* volume 8, pages 587–596.
- LaConti JJ, Laiakis EC, Mays AD, Peran I, Kim SE, Shay JW, Riegel AT, Fornace AJ Jr, Wellstein A. (2015) Distinct serum metabolomics profiles associated with malignant progression in the KrasG12D mouse model of pancreatic ductal adenocarcinoma. *BMC Genomics*; 16 Suppl 1: S1.
- Ladenson RP, Schwartz SO, Ivy AC. (1949) Incidence of the blood groups and the secretor factor in patients with pernicious anemia and stomach carcinoma. *Am. J. Med. Sci.* 217, 194–197.
- Lai X, Wang M, McElyea SD, Sherman S, House M, Korc M. (2017) A microRNA signature in circulating exosomes is superior to exosomal glypican-1 levels for diagnosing pancreatic cancer. *Cancer Lett*; 393: 86-93.
- Le Roith D, Bondy C, Yakar S, Liu JL, Butler A. (2001) 'The somatomedin hypothesis: 2001. *Endocr Rev.*; 22: 53-74.
- Leary RJ, Kinde I, Diehl F, Schmidt K, Clouser C, Duncan C, Antipova A, Lee C, Mckernan K, De La Vega FM, Kinzler KW, Vogelstein B, Diaz LA, Velculescu VE. (2010) Development of Personalized Tumor Biomarkers Using Massively Parallel Sequencing. *Science Translational Medicine. Sci Transl Med*; 2(20):20ra14
- Leask A. (2009) Death of a tumor: targeting CCN in pancreatic cancer. *J Cell Commun Signal* 3: 159 – 160
- Lee EJ, Gusev Y, Jiang J, Nuovo GJ, Lerner MR, Frankel WL, Morgan DL, Postier RG, Brackett DJ, Schmittgen TD. Expression profiling identifies microRNA signature in pancreatic cancer. *Int J Cancer* 2007; 120: 1046-1054.

Lee ES, Lee JM. (2014) Imaging diagnosis of pancreatic cancer: a state-of-the-art review. *World J Gastroenterol*; 20: 7864-7877. PMID: 24976723 DOI: 10.3748/wjg.v20.i24.7864

Lennon A.M, Wolfgang CL, Canto MI, Klein AP, Herman JM, Goggins M, Fishman EK, Kamel I, Weiss MJ, Diaz LA, Papadopoulos N, Kinzler KW, Vogelstein B, Hruban RH. (2014) The early detection of pancreatic cancer: What will it take to diagnose and treat curable pancreatic neoplasia? *Cancer Res.*, 74, 3381–33819.doi: 10.1158/0008-5472.CAN-14-0734

Li A, Yu J, Kim H, Wolfgang CL, Canto MI, Hruban RH, Goggins M. (2013) MicroRNA array analysis finds elevated serum miR-1290 accurately distinguishes patients with low-stage pancreatic cancer from healthy and disease controls. *Clin Cancer Res*; 19: 3600- 3610.

Li C, Simeone DM, Brenner DE, Anderson MA, Shedden KA, Ruffin MT, Lubman DM. (2009) Pancreatic cancer serum detection using a lectin/glyco-antibody array method. *J Proteome Res*; 8: 483-492. PMID: 19072160 DOI: 10.1021/pr8007013

Li D, Mallory T, Satomura S. (2001) AFP-L3: a new generation of tumor marker for hepatocellular carcinoma. *Clin Chim Acta*; 313: 15-19. PMID: 11694234 DOI: 10.1016/S0009-8981(01)00644-1

Li X, Ma Q, Xu Q, Duan W, Lei J, Wu E. (2012) Targeting the cancer-stroma interaction: a potential approach for pancreatic cancer treatment. *Curr Pharm Des.*;18(17):2404–15.

Liffers ST, Munding JB, Vogt M, Kuhlmann JD, Verdoodt B, Nambiar S, Maghnouj A, Mirmohammadsadegh A, Hahn SA, Tannapfel A. (2011) MicroRNA- 148a is down-regulated in human pancreatic ductal adenocarcinomas and regulates cell survival by targeting CDC25B. *Lab Invest*;91:1472-9. doi: 10.1038/labinvest.2011.99.

Liggett T, Melnikov A, Yi QL, Replogle C, Brand R, Kaul K, Talamonti M, Abrams RA, Levenson V. (2010) Differential methylation of cell-free circulating DNA among patients with pancreatic cancer versus chronic pancreatitis. *Cancer*; 116: 1674-1680. doi: 10.1002/cncr.24893.

Llop E, Ferrer-Batallé M, Barrabés S, Guerrero PE, Ramírez M, Saldoval R, Rudd PM, Aleixandre RN, Comet J, de Llorens R, Peracaula R. (2016) Improvement of Prostate Cancer Diagnosis by Detecting PSA Glycosylation-Specific Changes. *Theranostics*; 6: 1190-1204. PMID: 27279911 DOI: 10.7150/thno.15226

- Llop E, Guerrero P, Duran A, Barrabés S, Massaguer A, José Ferri M, Albiol-Quer M, de Llorens R, Peracaula R. (2018) Glycoprotein biomarkers for the detection of pancreatic ductal adenocarcinoma. *World J Gastroenterol*;24(24):2537-2554.
- Lo Cicero A, Stahl PD, Raposo G. (2015) Extracellular vesicles shuffling intercellular messages: for good or for bad. *Curr Opin Cell Biol*; 35: 69-77.
- Locker GY, Hamilton S, Harris J, Jessup JM, Kemeny N, Macdonald JS, Somerfield MR, Hayes DF, Bast RC Jr; ASCO. (2006) ASCO 2006 up- date of recommendations for the use of tumor markers in gastrointestinal cancer. *J Clin Oncol*; 24: 5313-5327.
- Loffreda S, Yang SQ, Lin HZ, Karp CL, Brengman ML, Wang DJ, Klein AS, Bulkley GB, Bao C, Noble PW, Lane MD, Diehl AM. (1998) Leptin regulates proinflammatory immune responses. *Faseb J*; 12: 57-65.
- Lowenfels AB, Maisonneuve P. (2006) Epidemiology and risk factors for pancreatic cancer. *Best Pract Res Clin Gastroenterol*, 20:197–209.
- Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, Sweet-Cordero A, Ebert BL, Mak RH, Ferrando AA, Downing JR, Jacks T, Horvitz HR, Golub TR. (2005) MicroRNA expression profiles classify human cancers. *Nature*; 435: 834-838.
- Lu X, Fang Y, Wang Z, Xie J, Zhan Q, Deng X, Chen H, Jin J, Peng C, Li H, Shen B. (2013) Downregulation of gas5 increases pancreatic cancer cell proliferation by regulating CDK6. *Cell Tissue Res*; 354: 891-896.
- Luo G, Liu C, Guo M, Cheng H, Lu Y, Jin K, Liu L, Long J, Xu J, Lu R, Ni Q, Yu X. (2017) Potential biomarkers in lewis negative patients with pancreatic cancer. *Ann Surg*; 265: 800-805.
- Lynge E. (1989) Screening for cancer of the cervix uteri. *World J Surg.*; 13:71–78.
- Madhavan B, Yue S, Galli U, Rana S, Gross W, Müller M, Giese NA, Kalthoff H, Becker T, Büchler MW, Zöller M. Combined evaluation of a panel of protein and miRNA serum-exosome biomarkers for pancreatic cancer diagnosis increases sensitivity and specificity. *Int J Cancer* 2015; 136: 2616-2627.
- Maire F, Micard S, Hammel P, Voitot H, Levy P, Cugnenc PH, Ruszniewski P, Puig PL. (2002) Differential diagnosis between chronic pancreatitis and pancreatic cancer: value of the detection of KRAS2 mutations in circulating DNA. *Br J Cancer*; 87: 551-554.
- Maitra A, Adsay NV, Argani P, Iacobuzio-Dona- hue C, De Marzo A, Cameron JL, Yeo CJ, Hruban RH. (2003) Multicomponent analysis of the pancreatic adenocarcinoma

progression model using a pancreatic intraepithelial neoplasia tissue microarray. *Mod Pathol*; 16: 902- 912.

Makawita S, Diamandis EP. (2010) The bottleneck in the cancer biomarker pipeline and protein quantification through mass spectrometry-based approaches: current strategies for candidate verification. *Clin Chem* 56(2):212–222.

Makawita S, Dimitromanolakis A, Soosaipillai A, Soleas I, Chan A, Gallinger S, Haun RS, Blasutig IM, Diamandis EP. (2013) Validation of four candidate pancreatic cancer serological biomarkers that improve the performance of CA19.9. *BMC Cancer*; 13: 404.

Mancera-Arteu M, Giménez E, Barbosa J, Peracaula R, Sanz-Nebot V. (2017) Zwitterionic-hydrophilic interaction capillary liquid chromatography coupled to tandem mass spectrometry for the characterization of human alpha-acid-glycoprotein N-glycan isomers. *Anal Chim Acta*; 991: 76-88. PMID: 29031301 DOI: 10.1016/j.aca.2017.07.068

Mandel JS, Church TR, Ederer F, Bond JH. (1999) Colorectal cancer mortality: effectiveness of biennial screening for fecal occult blood. *J Natl Cancer Inst.*; 91:434–437.

Marcos, N. T. Pinho S, Grandela C, Cruz A, Samyn-Petit B, Harduin-Lepers A, Almeida R, Silva F, Costa J, Kihlberg J, Clausen H, Reis CA. (2004) Role of the human ST6GalNAc-I and ST6GalNAc-II in the synthesis of the cancer- associated sialyl-Tn antigen. *Cancer Res.* 64, 7050–7057.

Marrelli D, Caruso S, Pedrazzani C, Neri A, Fernandes E, Marini M, Pinto E, Roviello F. (2009) CA19-9 serum levels in obstructive jaundice: clinical value in benign and malignant conditions. *Am J Surg*; 198: 333-339. PMID: 19375064 DOI: 10.1016/j.amjsurg.2008.12.031

Masamune A, Shimosegawa T. (2015) Pancreatic stellate cells: a dynamic player of the intercellular communication in pancreatic cancer. *Clin Res Hepatol Gastroenterol.*;39(Suppl 1): S98–103.

Matsubayashi H, Sato N, Fukushima N, Yeo CJ, Walter KM, Sahin F, Hruban RH, Goggins M. (2003) Methylation of cyclin D2 is observed frequently in pancreatic cancer but is also an age-related phenomenon in gastrointestinal tissues. *Clin Cancer Res*;9:1446-52.

Matsuda A, Higashi M, Nakagawa T, Yokoyama S, Kuno A, Yonezawa S, Narimatsu H. (2017) Assessment of tumor characteristics based on glycoform analysis of



membrane-tethered MUC1. *Lab Invest*; 97: 1103-1113. PMID: 28581490 DOI: 10.1038/labinvest.2017.53

Matsumoto H, Shinzaki S, Narisada M, Kawamoto S, Kuwamoto K, Moriwaki K, Kanke F, Satomura S, Kumada T, Miyoshi E. (2010) Clinical application of a lectin-antibody ELISA to measure fucosylated haptoglobin in sera of patients with pancreatic cancer. *Clin Chem Lab Med*; 48: 505-512. PMID: 20128732 DOI: 10.1515/CCLM.2010.095

Matsumoto I, Shirakawa S, Shinzeki M, Asari S, Goto T, Ajiki T, Fukumoto T, Kitajima K, Ku Y. (2013) 18-Fluorodeoxyglucose positron emission tomography does not aid in diagnosis of pancreatic ductal adenocarcinoma. *Clin Gastroenterol Hepatol*; 11: 712-718. PMID: 23353642 DOI: 10.1016/j.cgh.2012.12.033

Matsuno S, Egawa S, Fukuyama S., Motoi F, Sunamura M, Isaji S, Imaizumi T, Okada S, Kato H, Suda K, Nakao A, Hiraoka T, Hosotani R, Takeda K. (2004) Pancreatic Cancer Registry in Japan: 20 years of experience. *Pancreas*, 28, 219–230.

Mayers JR, Wu C, Clish CB, Kraft P, Torrence ME, Fiske BP, Yuan C, Bao Y, Townsend MK, Twofer SS, Davidson SM, Papagiannakopoulos T, Yang A, Dayton TL, Ogino S, Stampfer MJ, Giocannucci EL, Qian ZR, Robinson DA, Ma J, Sesso HD, Gaziano JM, Cochrane BB, Liu S, Wactawski-Wende J, Manson JE, Pollak MN, Kimmelman AC, Souza A, Pierce K, Wang TJ, Gerszten RE, Fuchs CS, Vander Heiden MG, Wolpin BM. (2014) Elevation of circulating branched-chain amino acids is an early event in human pancreatic adenocarcinoma development. *Nat Med.*; 20:1193–1198. doi: 10.1038/nm.3686.

McCoy Jr JP, Varani J, Goldstein IJ. (1983) Enzyme-linked lectin assay (ELLA): use of alkaline phosphatase-conjugated *Griffonia simplicifolia* B4 isolectin for the detection of alpha-D-galactopyranosyl end groups. *Analytical Biochemistry* 130(2):437–444. DOI 10.1016/0003-2697(83)90613-9.

McTiernan A. (2008) Mechanisms linking physical activity with cancer. *Nat Rev Cancer*; 8: 205–11.

Meany DL, Chan DW. (2011) Aberrant glycosylation associated with enzymes as cancer biomarkers. *Clin Proteomics*; 8: 7. PMID: 21906357 DOI: 10.1186/1559-0275-8-7

Melnikov AA, Scholtens D, Talamonti MS, Bentrem DJ, Levenson VV. (2009) Methylation profile of circulating plasma DNA in patients with pancreatic cancer. *J Surg Oncol*; 99: 119- 122.

Melo SA, Luecke LB, Kahlert C, Fernandez AF, Gammon ST, Kaye J, LeBleu VS, Mittendorf EA, Weitz J, Rahbari N, Reisfelder C, Pilarsky C, Fraga MF, Piwnica-Worms D, Kalluri R. (2015) Glypican-1 identifies cancer exosomes and detects early pancreatic cancer. *Nature*; 523: 177-82. doi: 10.1038/nature14581.

Mody, S, Joshi, W, Chaney, J. (1995) *Pharmacol. Toxicol. Methods* 33;1–10.

Mordente A, Meucci E, Martorana G, Silvestrini A. (2015) *Cancer Biomarkers Discovery and Validation: State of the Art, Problems and Future Perspectives*, Springer Science + Business Media Dordrecht 9, R. Scatena (ed.), *Advances in Cancer Biomarkers, Advances in Experimental Medicine and Biology* 867.

Moremen KW, Tiemeyer M, Nairn, A. V. (2012) Vertebrate protein glycosylation: diversity, synthesis and function. *Nat. Rev. Mol. Cell Biol.* 13, 448–462.

Mulcahy HE, Lyautey J, Lederrey C, qi Chen X, Anker P, Alstead EM, Ballinger A, Farthing MJ, Stroun M. (1998) A prospective study of K-ras mutations in the plasma of pancreatic cancer patients. *Clin Cancer Res*; 4: 271-275.

Muller S, Raulefs S, Bruns P, Afonso-Grunz F, Plotner A, Thermann R, Jager C, Schlitter AM, Kong B, Regel I, Roth WK, Rotter B, Hoffmeier K, Kahl G, Koch I, Theis FJ, Kleeff J, Winter P, Michalski CW. (2015) Next-generation sequencing reveals novel differentially regulated mRNAs, lncRNAs, miRNAs, sdRNAs and a piR-NA in pancreatic cancer. *Mol Cancer*; 14: 94.

Munkley J, Elliott DJ. (2016) Hallmarks of glycosylation in cancer. *Oncotarget*; 7: 35478-35489. PMID: 27007155 DOI: 10.18632/oncotarget.8155

Nagrath S, Sequist LV, Maheswaran S, Bell DW, Irimia D, Ulkus L, Smith MR, Kwak EL, Digumarthy S, Muzikansky A, Ryan P, Balis UJ, Tompkins RG, Haber DA, Toner M. (2007) Isolation of rare circulating tumor cells in cancer patients by microchip technology. *Nature*; 450: 1235-1239.

Nakajima K, Inomata M, Iha H, Hiratsuka T, Etoh T, Shiraishi N, Kashima K, Kitano S. (2015) Establishment of new predictive markers for distant recurrence of colorectal cancer using lectin microarray analysis. *Cancer Medicine* 4(2):293–302. DOI 10.1002/cam4.342.

Nakamori, S, Kameyama M, Imaoka S, Furukawa H, Ishikawa O, Sasaki Y, Kabuto T, Iwanaga T, Matsushita Y, Irimura T. (1993) Increased expression of sialyl Lewis<sup>x</sup> antigen correlates with poor survival in patients with colorectal carcinoma: clinicopathological and immunohistochemical study. *Cancer Res.* 53, 3632–3637.

Nakata D. (2017) Increased N-glycosylation of Asn<sup>88</sup> in serum pancreatic ribonuclease 1 is a novel diagnostic marker for pancreatic cancer. *Sci Rep* 2014; 4: 6715, PMID: 25336120 DOI: 10.1038/srep06715

Nakchbandi I, Lang R, Kinder B, Insogna K. (2008). The Role of the Receptor Activator of Nuclear Factor- $\kappa$ B Ligand/Osteoprotegerin Cytokine System in Primary Hyperparathyroidism. *The Journal of clinical endocrinology and metabolism*. 93. 967-73. 10.1210/jc.2007-1645. DOI: 10.1210/jc.2007-1645

National Cancer Institute (2018) Lifetime risk tables. [https://seer.cancer.gov/csr/1975\\_2015/results\\_merged/topic\\_lifetime\\_risk.pdf](https://seer.cancer.gov/csr/1975_2015/results_merged/topic_lifetime_risk.pdf)  
Accessed 31 Jan. 2019

National Cancer Registration and Analysis Service. Stage Breakdown by CCG 2014. London: NCRAS; 2016.

National Comprehensive Cancer Network. 2017 [accessed, February 2019]. In: NCCN guidelines version 3.2017: pancreatic adenocarcinoma [Internet]. Available from: URL: [https://www.nccn.org/professionals/physician\\_gls/pdf/pancreatic.pdf](https://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf)

Newman AM, Bratman SV, To J, Wynne JF, Eclov NC, Modlin LA, Liu CL, Neal JW, Wakelee HA, Merritt RE, Shrager JB, Loo BW Jr, Alizadeh AA, Diehn M. (2014) An ultrasensitive method for quantitating circulating tumor DNA with broad patient coverage. *Nat Med*; 20: 548- 554.

Ni XG, Bai XF, Mao YL, Shao YF, Wu JX, Shan Y, Wang CF, Wang J, Tian YT, Liu Q, Xu DK, Zhao P. (2005) The clinical value of serum CEA, CA19-9, and CA242 in the diagnosis and prognosis of pancreatic cancer. *Eur J Surg Oncol*; 31: 164-169. PMID: 15698733 DOI: 10.1016/j.ejso.2004.09.007

Nie S, Lo A, Wu J, Zhu J, Tan Z, Simeone DM, Anderson MA, Shedden KA, Ruffin MT, Lubman DM. (2014) Glycoprotein biomarker panel for pancreatic cancer discovered by quantitative proteomics analysis. *J Proteome Res*; 13: 1873-1884. PMID: 24571389 DOI: 10.1021/pr400967x

Nyström L, Andersson I, Bjurstam N, Frisell J, Nordenskjöld B, Rutqvist. (2002) Long-term effects of mammography screening: updated overview of the Swedish randomised trials. *Lancet*; 359:909–919.

O'Brien DP, Sandanayake NS, Jenkinson C, Gentry-Maharaj A, Apostolidou S, Fourkala EO, Camuzeaux S, Blyuss O, Gunu R, Dawnay A, Zaikin A, Smith RC, Jacobs IJ, Menon U, Costello E, Pereira SP, Timms JF. (2015) Serum CA19-9 is significantly upregulated up to 2 years before diagnosis with pancreatic cancer:

implications for early disease detection. Clin Cancer Res; 21: 622-631. PMID: 24938522 DOI: 10.1158/1078-0432.CCR-14-0365

O'Flaherty JD, Gray S, Richard D, Fennell D, O'Leary JJ, Blackhall FH, O'Byrne KJ. (2012) Circulating tumor cells, their role in metastasis and their clinical utility in lung cancer. Lung Cancer; 76: 19-25.

O'Neill CB, Atoria CL, O'Reilly EM, LaFemina J, Henman MC, Elkin EB. (2012) Costs and trends in pancreatic cancer treatment. Cancer; 118: 5132-5139. PMID: 22415469 DOI: 10.1002/cncr.27490

Ozdemir BC, Pentcheva-Hoang T, Carstens JL, Zheng X, Wu CC, Simpson TR, Laklai H, Sugimoto H, Kahlert C, Novitskiy SV, De Jesus-Acosta A, Sharma P, Heidari P, Mahmood U, Chin L, Moses HL, Weaver VM, Maitra A, Allison JP, LeBleu VS, Kalluri R. (2014) Depletion of carcinoma-associated fibroblasts and fibrosis induces immunosuppression and accelerates pancreas cancer with reduced survival. Cancer Cell; 25: 719-734.

Pan S, Chen R, Crispin DA, May D, Stevens T, McIntosh MW, Bronner MP, Ziogas A, Anton-Culver H, Brentnall TA. (2011) Protein alterations associated with pancreatic cancer and chronic pancreatitis found in human plasma using global quantitative proteomics profiling. J Proteome Res 10: 2359 – 2376.

Pan S, Chen R, Tamura Y, Crispin DA, Lai LA, May DH, McIntosh MW, Goodlett DR, Brentnall TA. (2014) Quantitative glycoproteomics analysis reveals changes in N-glycosylation level associated with pancreatic ductal adenocarcinoma. J Proteome Res; 13: 1293-1306. PMID: 24471499 DOI: 10.1021/pr4010184

Pantel K, Brakenhoff RH. (2004) Dissecting the metastatic cascade. Nat Rev Cancer; 4:448-456.

Paredes, J, Figueiredo J, Albergaria A, Oliveira P, Carvalho J, Ribeiro AS, Caldeira J, Costa AM, Simões-Correia J, Oliveira MJ, Pinheiro H, Pinho SS, Mateus R, Reis CA, Leite M, Fernandes MS, Schmitt F, Carneiro F, Figueiredo C, Oliveira C, Seruca R. (2012) Epithelial E- and P-cadherins: role and clinical significance in cancer. Biochim. Biophys. Acta 1826, 297–311. doi: 10.1016/j.bbcan.2012.05.002.

Park HD, Kang ES, Kim JW, Lee KT, Lee KH, Park YS, Park JO, Lee J, Heo JS, Choi SH, Choi DW, Kim S, Lee JK, Lee SY. (2012) Serum CA19-9, cathepsin D, and matrix metalloproteinase-7 as a diagnostic panel for pancreatic ductal adenocarcinoma. Proteomics; 12: 3590- 3597.

Park R, Kim M, So H, Jung B, Moon S, Chung S, Ko C, Kim B, Chung H. (2000) *Biochem. Pharmacol.* 60; 1685–1691.

Parkin DM, Boyd L, Walker LC. (2010) The fraction of cancer attributable to lifestyle and environmental factors in the UK in 2010. *Br J Cancer* 2011; 105 Suppl 2: S77-S81.

Pavlou MP, Diamandis EP, Blasutig IM. (2013) The long journey of cancer biomarkers from the bench to the clinic. *Clin Chem*; 59: 147-157. PMID: 23019307 DOI: 10.1373/clinchem.2012.184614

Peng W, Gao W, Feng J. (2014) Long noncoding RNA HULC is a novel biomarker of poor prognosis in patients with pancreatic cancer. *Med Oncol*; 31: 346.

Peracaula R, Barrabés S, Sarrats A, Rudd PM, de Llorens R. (2008) Altered glycosylation in tumours focused to cancer diagnosis. *Dis Markers*; 25: 207-218. PMID: 19126965 DOI: 10.1155/2008/797629

Pérez-Garay M, Arteta B, Llop E, Cobler L, Pagès L, Ortiz R, Ferri MJ, de Bolós C, Figueras J, de Llorens R, Vidal-Vanaclocha F, Peracaula R. (2013)  $\alpha$ 2,3-Sialyltransferase ST3Gal IV promotes migration and metastasis in pancreatic adenocarcinoma cells and tends to be highly expressed in pancreatic adenocarcinoma tissues. *Int J Biochem Cell Biol*; 45: 1748-1757. PMID: 23726834 DOI: 10.1016/j.biocel.2013.05.015

Peumans WJ, Van Damme JM, Barre A, Rougé P. (2001) Classification of Plant Lectins in families of structurally and evolutionary related proteins. In: *The molecular immunology of complex carbohydrates -2*. Boston: Springer, 27–54.

Pinho SS, Carvalho S, Marcos-Pinto R, Magalhães A, Oliveira C, Gu J, Dinis-Ribeiro M, Carneiro F, Seruca R, Reis CA. (2013) Gastric cancer: adding glycosylation to the equation. *Trends Mol. Med.* 19, 664–676. doi: 10.1016/j.molmed.2013.07.003.

Pinho SS, Oliveira P, Cabral J, Carvalho S, Huntsman D, Gärtner F, Seruca R, Reis CA, Oliveira C. (2012) Loss and recovery of Mgat3 and GnT-III Mediated E-cadherin N-glycosylation is a mechanism involved in epithelial-mesenchymal- epithelial transitions. *PLoS ONE* 7, e33191. doi: 10.1371/journal.pone.0033191. Epub 2012 Mar 13.

Pinho SS, Reis CA. (2015) Glycosylation in cancer: mechanisms and clinical implications. *Nat Rev Cancer*; 15: 540-555. PMID: 26289314 DOI: 10.1038/nrc3982

Pinho SS, Seruca R, Gärtner F, Yamaguchi Y, Gu J, Taniguchi N, Reis CA. (2011) Modulation of E-cadherin function and dysfunction by N-glycosylation. *Cell. Mol. Life Sci.* 68, 1011–1020. doi: 10.1007/s00018-010-0595-0.

Pishvaian MJ, Joseph Bender R, Matrisian LM, Rahib L, Hendifar A, Hoos WA, Mikhail S, Chung V, Picozzi V, Heartwell C, Mason K, Varieur K, Aberra M, Madhavan S, Petricoin E, Brody JR. (2016) A pilot study evaluating concordance between blood-based and patient-matched tumor molecular testing within pancreatic cancer patients participating in the Know Your Tumor (KYT) initiative. *Oncotarget*; 8: 83446-83456. PMID: 29137355 DOI: 10.18632/oncotarget.13225

Pistol-Tanase C, Raducan E, Dima SO, Albulescu L, Alina I, Marius P, Cruceru LM, Codorean E, Neagu TM and Popescu I. (2008) Assessment of soluble angiogenic markers in pancreatic cancer. *Biomark Med*; 2: 447-455.

Polanski M, Anderson NL. (2007) A list of candidate cancer biomarkers for targeted proteomics. *Biomark Insights*; 1: 1-48. PMID: 19690635 DOI: 10.1177/117727190600100001

Poruk KE, Firpo MA, Adler DG, Mulvihill SJ. (2013) Screening for pancreatic cancer: why, how, and who? *Ann Surg*; 257: 17-26. PMID: 22895395 DOI: 10.1097/SLA.0b013e31825ffbfbb

Zhou B, Xu JW, Cheng YG, Gao JY, Hu SY, Wang L, Zhan HX. (2017) Early detection of pancreatic cancer: Where are we now and where are we going? *Int J Cancer*; 141: 231-241. PMID: 28240774 DOI: 10.1002/ijc.30670

Pothula SP, Xu Z, Goldstein D, Pirola RC, Wilson JS, Apte MV. (2016) Key role of pancreatic stellate cells in pancreatic cancer. *Cancer Lett.*;381(1):194- 200.

Pour PM, Tempero MM, Takasaki H, Uchida E, Takiyama Y, Burnett DA, Steplewski Z. (1988) Expression of blood group-related antigens ABH, Lewis A, Lewis B, Lewis X, Lewis Y, and CA 19-9 in pancreatic cancer cells in comparison with the patient's blood group type. *Cancer Res*; 48: 5422-5426. PMID: 3166398

Puli SR, Bechtold ML, Buxbaum JL, Eloubeidi MA. (2013) How good is endoscopic ultrasound-guided fine-needle aspiration in diagnosing the correct etiology for a solid pancreatic mass?: A meta-analysis and systematic review. *Pancreas*; 42: 20-26 .

Quinn JJ, Chang HY. (2016) Unique features of long non-coding RNA biogenesis and function. *Nat Rev Genet*; 17: 47-62.

Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. (2014) Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. *Cancer Res*; 74: 2913-2921. PMID: 24840647 DOI: 10.1158/0008-5472.CAN-14-0155

Reis C, Osorio H, Silva L, Gomes C, David L. (2010) Alterations in glycosylation as biomarkers for cancer detection. *J. Clin. Pathol.* 63, 322–329.

Remmers N, Anderson JM, Linde EM, DiMaio DJ, Lazenby AJ, Wandall HH, Mandel U, Clausen H, Yu F, Hollingsworth MA. (2013) Aberrant expression of mucin core proteins and o-linked glycans associated with progression of pancreatic cancer. *Clin Cancer Res*; 19: 1981-1993. PMID: 23446997 DOI: 10.1158/1078-0432.CCR-12-2662

Ren S, Wang F, Shen J, Sun Y, Xu W, Lu J, Wei M, Xu C, Wu C, Zhang Z, Gao X, Liu Z, Hou J, Huang J and Sun Y. (2013) Long non-coding RNA metastasis associated in lung adenocarcinoma transcript 1 derived miniRNA as a novel plasma-based biomarker for diagnosing prostate cancer. *Eur J Cancer*; 49: 2949-2959.

Resovi A, Bani M, Porcu L, Anastasia A, Minoli L, Allavena P, Cappello P, Novelli F, Scarpa A, Morandi E, Falanga A, Torri V, Taraboletti G, Belotti D, Giavazzi R. (2018) Soluble stroma-related biomarkers of pancreatic cancer, *EMBO Mol Med* 10: e8741.

Rexrode KM, Pradhan A, Manson JE et al... (2003) 'Relationship of total and abdominal adiposity with CRP and IL-6 in women. *Ann Epidemiol*; 13: 674-82.

Rhim AD, Mirek ET, Aiello NM, Maitra A, Bailey JM, McAllister F, Reichert M, Beatty GL, Rustgi AK, Vonderheide RH, Leach SD, Stanger BZ. (2012) EMT and dissemination precede pancreatic tumor formation. *Cell*; 148: 349-361.

Rhim AD, Oberstein PE, Thomas DH, Mirek ET, Palermo CF, Sastra SA, Dekleva EN, Saunders T, Becerra CP, Tattersall IW, Westphalen CB, Kitajewski J, Fernandez-Barrena MG, Fernandez-Zapico ME, Iacobuzio-Donahue C, Olive KP and Stanger BZ. (2014) Stromal elements act to restrain, rather than support, pancreatic ductal adenocarcinoma. *Cancer Cell*; 25: 735- 747.

Rhim AD, Thege FI, Santana SM, Lannin TB, Saha TN, Tsai S, Maggs LR, Kochman ML, Ginsberg GG, Lieb JG, Chandrasekhara, Ahmad N, Yang YX, Kirby BJ, Stanger BZ. (2014) Detection of circulating pancreas epithelial cells in patients with pancreatic cystic lesions. *Gastroenterology*; 146:647-51. doi: 10.1053/j.gastro.2013.12.007

Ribeiro A, Monteiro S, Carrapiço B, Ferreira R. (2014) *Molecules* 19; 20350–20373.

Ritchie SA, Akita H, Takemasa I, Eguchi H, Pastural E, Nagano H, Monden M, Doki Y, Mori M, Jin W, Sajobi TT, Jayasinghe D, Chitou B, Yamazaki Y, White T and Goodenowe DB. (2013) Metabolic system alterations in pancreatic cancer patient serum: potential for early detection. *BMC Cancer*; 13: 416.

Riva F, Dronov OI, Khomenko DI, Huguet F, Louvet C, Mariani P, Stern MH, Lantz O, Proudhon C, Pierga JY, Bidard FC. (2016) Clinical applications of circulating tumor DNA and circulating tumor cells in pancreatic cancer. *Mol Oncol*; 10: 481-493. PMID: 26856794 DOI: 10.1016/j.molonc.2016.01.006

Robbins PD, Morelli AE. (2014) Regulation of immune responses by extracellular vesicles. *Nat Rev Immunol*; 14: 195-208.

Melo SA, Sugimoto H, O'Connell JT, Kato N, Villanueva A, Vidal A, Qiu L, Vitkin E, Perelman LT, Melo CA, Lucci A, Ivan C, Calin GA and Kalluri R. (2014) Cancer exosomes perform cell-independent microRNA biogenesis and promote tumorigenesis. *Cancer Cell*; 26: 707-721.

Roldo C, Missiaglia E, Hagan JP, Falconi M, Capelli P, Bersani S, Calin GA, Volinia S, Liu CG, Scarpa A and Croce CM. (2006) MicroRNA expression abnormalities in pancreatic endocrine and acinar tumors are associated with distinctive pathologic features and clinical behavior. *J Clin Oncol*; 24: 4677-4684. DOI: 10.3390/molecules19022458

Rong Y, Jin D, Hou C, Hu J, Wu W, Ni X, Wang D, Lou W. (2010) Proteomics analysis of serum protein profiling in pancreatic cancer patients by DIGE: up-regulation of mannose-binding lectin 2 and myosin light chain kinase 2. *BMC Gastroenterol* 10: 68.

Root A, Allen P, Tempst P, Yu K. (2018) Protein Biomarkers for Early Detection of Pancreatic Ductal Adenocarcinoma: Progress and Challenges, *Cancers* 2018, 10, 6.

Rosen SD, Bertozzi CR. (1994) The selectins and their ligands. *Curr. Opin. Cell Biol.* 6, 663–673.

Roth J. (2011) Lectins for histochemical demonstration of glycans. *Histochemistry and Cell Biology* 136(2):117–130. DOI 10.1007/s00418-011-0848-5.

Roth Z, Yehezkel G, Khalaila I. (2012) Identification and quantification of protein glycosylation. *International Journal of Carbohydrate Chemistry* 2012:640923. DOI 10.1155/2012/640923.

Rulyak SJ, Lowenfels AB, Maisonneuve P and Brentnall TA. (2003) Risk factors for the development of pancreatic cancer in familial pancreatic cancer kindreds. *Gastroenterology*; 124: 1292-1299.

Ryu JK, Hong SM, Karikari CA, Hruban RH, Goggins MG, Maitra A. (2010) Aberrant MicroRNA-155 expression is an early event in the multistep progression of pancreatic adenocarcinoma. *Pancreatol*; 10: 66-73.



Sakai A, Suzuki M, Kobayashi T, Nishiumi S, Yamanaka K, Hirata Y, Nakagawa T, Azuma T, Yoshida M. (2016) Pancreatic cancer screening using a multiplatform human serum metabolomics system. *Biomark Med*; 10: 577- 586. doi: 10.2217/bmm-2016-0020.

San Lucas FA, Allenson K, Bernard V, Castillo J, Kim DU, Ellis K, Ehli EA, Davies GE, Petersen JL, Li D, Wolff R, Katz M, Varadhachary G, Wistuba I, Maitra A, Alvarez H. (2016) Minimally invasive genomic and transcriptomic profiling of visceral cancers by next-generation sequencing of circulating exosomes. *Ann Oncol*; 27: 635-641.

Sarrats A, Comet J, Tabarés G, Ramírez M, Aleixandre RN, de Llorens R, Peracaula R. (2010) Differential percentage of serum prostate- specific antigen subforms suggests a new way to improve prostate cancer diagnosis. *Prostate*; 70: 1-9. PMID: 19670261 DOI: 10.1002/pros.21031

Sarrats A, Saldova R, Pla E, Fort E, Harvey DJ, Struwe WB, de Llorens R, Rudd PM, Peracaula R. (2010) Glycosylation of liver acute- phase proteins in pancreatic cancer and chronic pancreatitis. *Proteomics Clin Appl*; 4: 432-448. PMID: 21137062 DOI: 10.1002/prca.200900150

Sato N, Fukushima N, Maitra A, Matsubayashi H, Yeo CJ, Cameron JL, Hruban RH, Goggins M. (2003) Discovery of novel targets for aberrant methylation in pancreatic carcinoma using high-throughput microarrays. *Cancer Res*; 63: 3735-3742.

Sawey ET, Johnson JA, Crawford HC. (2007) Matrix metalloproteinase 7 controls pancreatic acinar cell transdifferentiation by activating the Notch signaling pathway. *Proc Natl Acad Sci USA* 104: 19327 – 19332.

Scaiewicz V, Sorin V, Fellig Y, Birman T, Mizrahi A, Galula J, Abu-Lail R, Shneider T, Ohana P, Buscail L, Hochberg A and Czerniak A. (2010) Use of H19 gene regulatory sequences in DNA-based therapy for pancreatic cancer. *J Oncol*; 2010: 178174. doi: 10.1155/2010/178174.

Schiess R, Wollscheid B, Aebersold R. (2009) Targeted proteomic strategy for clinical biomarker discovery. *Mol Oncol*; 3: 33-44. PMID: 19383365 DOI: 10.1016/j.molonc.2008.12.001

Schietinger A, Philip M, Yoshida BA, Azadi P, Liu H, Meredith SC, Schreiber H. (2006) A mutant chaperone converts a wild-type protein into a tumor-specific antigen. *Science* 314, 304–308.

Schoen RE, Pinsky PF, Weissfeld JL, Yokochi LA, Church T, Laiyemo AO, Bresalier R, Andriole GL, Buys SS, Crawford ED, Fouad MN, Isaacs CC, Reding DJ, O'Brien B,

Carrick DM, Wright P, Riley TL, Purdue MP, Izmirlian G, Kramer BS, Miller AB, Gohagan JK, Prorok PC, Berg CD, PLCO Project Team. (2012) Colorectal-cancer incidence and mortality with screening flexible sigmoidoscopy. *N Engl J Med*. 2012; 366:2345–2357. doi: 10.1056/NEJMoa1114635.

Schultz NA, Dehlendorff C, Jensen BV, Bjerregaard JK, Nielsen KR, Bojesen SE, Calatayud D, Nielsen SE, Yilmaz M, Hollander NH, Andersen KK, Johansen JS. (2014) MicroRNA biomarkers in whole blood for detection of pancreatic cancer. *JAMA*; 311: 392-404.

Schumacher TN, Schreiber RD. (2015) Neoantigens in cancer immunotherapy. *Science*; 348: 69-74.

Secretan B, Straif K, Baan R, Grosse Y, El Ghissassi F, Bouvard V, Benbrahim-Tallaa L, Guha N, Freeman C, Galichet L, Coglian V, WHO International Agency for Research on Cancer Monograph Working Group. (2009). A review of human carcinogens--Part E: tobacco, areca nut, alcohol, coal smoke, and salted fish. *Lancet Oncol*,; 10: 1033-4.

Sefrioui D, Blanchard F, Toure E, Basile P, Beaussire L, Dolfus C, Perdrix A, Paresy M, Antonietti M, Iwanicki-Caron I, Alhameedi R, Lecleire S, Gangloff A, Schwarz L, Clatot F, Tuech JJ, Frébourg T, Jardin F, Sabourin JC, Sarafan-Vasseur N, Michel P, Di Fiore F. (2017) Diagnostic value of CA19.9, circulating tumour DNA and circulating tumour cells in patients with solid pancreatic tumours. *Br J Cancer*; 117: 1017-1025. PMID: 28772284 DOI: 10.1038/ bjc.2017.250

Shan S, Tanaka H, Shoyama Y. (2001) Enzyme-linked immunosorbent assay for glycyrrhizin using anti-glycyrrhizin monoclonal antibody and an eastern blotting technique for glucuronides of glycyrrhetic acid. *Analytical Chemistry* 73(24):5784–5790. DOI 10.1021/ac0106997.

Sharon N, Lis H. (2004) History of lectins: from hemagglutinins to biological recognition molecules. *Glycobiology* 14:53R–62R. DOI 10.1093/glycob/cwh122.

Shaw VE, Lane B, Jenkinson C, Cox T, Green- half W, Halloran CM, Tang J, Sutton R, Neoptolemos JP, Costello E. (2014) Serum cytokine biomarker panels for discriminating pancreatic cancer from benign pancreatic disease. *Mol Cancer*; 13: 114.

Shi C, Hruban RH, Klein AP. (2009) Familial pancreatic cancer. *Arch Pathol Lab Med*, 133:365–374.

Shi S, Yu X. (2018) Selecting chemotherapy for pancreatic cancer: Far away or so close?, Elsevier Inc, *Seminars in Oncology*, 0093-7754.

Siegel R, Miller K, Jema A. (2018), Cancer Statistics CA CANCER J CLIN 2018; 68:7–30.

Sigurdsson K. (1999) The Icelandic and Nordic cervical screening programs: trends in incidence and mortality rates through 1995. *Acta Obstet Gynecol Scand.*; 78:478–485.

Silva ML. (2015) Cancer serum biomarkers based on aberrant post- translational modifications of glycoproteins: Clinical value and discovery strategies. *Biochim Biophys Acta*; 1856: 165-177. PMID: 26232626 DOI: 10.1016/j.bbcan.2015.07.002

Singh S, Pal K, Yadav J, Tang H, Partyka K, Kletter D, Hsueh P, Ensink E, Kc B, Hostetter G, Xu HE, Bern M, Smith DF, Mehta AS, Brand R, Melcher K, Haab BB. (2015) Upregulation of glycans containing 3' fucose in a subset of pancreatic cancers uncovered using fusion- tagged lectins. *J Proteome Res*; 14: 2594-2605. PMID: 25938165 DOI: 10.1021/acs.jproteome.5b00142

Singh S, Tang SJ, Sreenarasimhaiah J, Lara LF, Siddiqui A. (2011) The clinical utility and limitations of serum carbohydrate antigen (CA19-9) as a diagnostic tool for pancreatic cancer and cholangiocarcinoma. *Dig Dis Sci*; 56: 2491-2496.

Skog J, Würdinger T, van Rijn S, Meiker DH, Gainche L, Sena-Esteves M, Curry WT Jr, Carter BS, Krichevsky AM, Breakefield XO. (2008) Glioblastoma microvesicles transport RNA and proteins that promote tumor growth and provide diagnostic biomarkers. *Nat Cell Biol*;10:1470-6. doi: 10.1038/ncb1800. Epub 2008 Nov 16.

Slater EP, Strauch K, Rospleszcz S, Ramaswamy A, Esposito I, Kloppel G, Matthai E, Heeger K, Fendrich V, Langer P, Bartsch DK. (2014) MicroRNA-196a and -196b as potential biomarkers for the early detection of familial pancreatic cancer. *Transl Oncol*; 7: 464-471.

Slebos RJ, Hoppin JA, Tolbert PE, Holly EA, Brock JW, Zhang RH, Bracci PM, Foley J, Stockton P, McGregor LM, Flake GP, Taylor JA. (2000) K-ras and p53 in pancreatic cancer: association with medical history, histopathology, and environmental exposures in a population-based study. *Cancer Epidemiol Biomarkers Prev*; 9: 1223-1232

Sobral AP, Rego MJ, Cavalcanti CL, Carvalho Jr LB, Beltrao EI. (2010) ConA and UEA-I lectin histochemistry of parotid gland mucoepidermoid carcinoma. *Journal of Oral Science* 52(1):49–54. DOI 10.2334/josnurd.52.49.

Sousa CM, Biancur DE, Wang X, Halbrook CJ, Sherman MH, Zhang L, Kremer D, Hwang RF, Witkiewicz AK, Ying H, Asara JM, Evans RM, Cantley LC, Lyssiotis CA, Kimmelman AC. (2016) Pancreatic stellate cells support tumor metabolism through autophagic alanine secretion. *Nature*; 536: 479-483.

- Sperti C, Pasquali C, Bissoli S, Chierichetti F, Liessi G, Pedrazzoli S. (2010) Tumor relapse after pancreatic cancer resection is detected earlier by 18-FDG PET than by CT. *J Gastrointest Surg*; 14: 131-140. PMID: 19777315 DOI: 10.1007/s11605-009-1010-8]
- St Laurent G, Wahlestedt C, Kapranov P. (2011) The Landscape of long noncoding RNA classification. *Trends Genet* 2015; 31: 239-251. Prensner JR and Chinnaiyan AM. The emergence of lncRNAs in cancer biology. *Cancer Discov*; 1: 391-407.
- Stathis A, Moore MJ. (2010) Advanced Pancreatic Carcinoma: Current Treatment and Future Challenges. *Nat Rev Clin Oncol*. 7(3):163.
- Stowell SR, Ju T, Cummings RD. (2015) Protein glycosylation in cancer. *Annu Rev Pathol*; 10: 473-510. PMID: 25621663 DOI: 10.1146/annurev-pathol-012414-040438]
- Szafranska AE, Davison TS, John J, Cannon T, Sipos B, Maghnouj A, Labourir E, Hahn SA. (2007) MicroRNA expression alterations are linked to tumorigenesis and non- neoplastic processes in pancreatic ductal adenocarcinoma. *Oncogene*;26:4442-52.
- Tabár L, Vitak B, Chen HH, Duffy SW, Yen MF, Chiang CF, Kruse UB, Tot T, Smith RA. (2000) The Swedish two-county trial twenty years later. Updated mortality results and new insights from long-term follow-up. *Radiol Clin N Am*; 38:625–651.
- Tabarés G, Radcliffe CM, Barrabés S, Ramírez M, Aleixandre RN, Hoesel W, Dwek RA, Rudd PM, Peracaula R, de Llorens R. (2006) Different glycan structures in prostate-specific antigen from prostate cancer sera in relation to seminal plasma PSA. *Glycobiology*; 16: 132-145. PMID: 16177264 DOI: 10.1093/glycob/cwj042
- Tahira AC, Kubrusly MS, Faria MF, Dazzani B, Fonseca RS, Maracaja-Coutinho V, Verjovski- Almeida S, Machado MC, Reis EM. (2011) Long noncoding intronic RNAs are differentially expressed in primary and metastatic pancreatic cancer. *Mol Cancer*; 10: 141.
- Takeuchi H, Haltiwanger RS. (2014) Significance of glycosylation in Notch signaling. *Biochem. Biophys. Res. Commun*. 453, 235–242.
- Tan Z, Yin H, Nie S, Lin Z, Zhu J, Ruffin MT, Anderson MA, Simeone DM, Lubman DM. (2015) Large-scale identification of core- fucosylated glycopeptide sites in pancreatic cancer serum using mass spectrometry. *J Proteome Res*; 14: 1968-1978. PMID: 25732060 DOI:10.1021/acs.jproteome.5b00068

Tang H, Partyka K, Hsueh P, Sinha JY, Kletter D, Zeh H, Huang Y, Brand RE, Haab BB. (2016) Glycans related to the CA19-9 antigen are elevated in distinct subsets of pancreatic cancers and improve diagnostic accuracy over CA19-9. *Cell Mol Gastroenterol Hepatol*; 2: 201-221.e15. PMID: 26998508 DOI: 10.1016/j.jcmgh.2015.12.003

Taniguchi N, Hancock W, Lubman D, Rudd P. (2009) The second golden age of glycomics: from functional glycomics to clinical applications. *J. Proteome Res.* 8, 425–426.

Terao N, Takamatsu S, Minehira T, Sobajima T, Nakayama K, Kamada Y, Miyoshi E. (2015) Fucosylation is a common glycosylation type in pancreatic cancer stem cell-like phenotypes. *World J Gastroenterol*; 21: 3876-3887. PMID: 25852272 DOI: 10.3748/wjg.v21.i13.3876

Tomaino B, Cappello P, Capello M, Fredolini C, Sperduti I, Migliorini P, Salacone P, Novarino A, Giacobino A, Ciuffreda L, Alessio M, Nistico P, Scarpa A, Pederzoli P, Zhou W, Petricoin Iii EF, Liotta LA, Giovarelli M, Milella M, Novelli F. (2011) Circulating autoantibodies to phosphorylated alpha-enolase are a hallmark of pancreatic cancer. *J Proteome Res*; 10: 105-112. doi: 10.1021/pr100213b.

Trajkovic K, Hsu C, Chiantia S, Rajendran L, Wenzel D, Wieland F, Schwille P, Brügger B, Simmons M. (2008) Ceramide triggers budding of exosome vesicles into multivesicular endosomes. *Science*;319:1244-7. doi: 10.1126/science.1153124.

Treadwell JR, Zafar HM, Mitchell MD, Tipton K, Teitelbaum U, Jue J. (2016) Imaging Tests for the Diagnosis and Staging of Pancreatic Adenocarcinoma: A Meta-Analysis. *Pancreas*; 45: 789-795.

Tseng J, Warshaw A, Sahani D, Lauwers G, Rattner D, Fernandez-del Castillo C. (2005) Serous Cystadenoma of the Pancreas Tumor Growth Rates and Recommendations for Treatment, *Ann Surg*;242: 413– 421.

Uemura T, Hibi K, Kaneko T, Takeda S, Inoue S, Okochi O, Nagasaka T, Nakao A. (2004) Detection of K-ras mutations in the plasma DNA of pancreatic cancer patients. *J Gastroenterol*; 39: 56-60.

Ulitsky I, Bartel DP. (2013) lincRNAs: genomics, evolution, and mechanisms. *Cell*; 154: 26-46.

Valadi H, Ekstrom K, Bossios A, Sjostrand M, Lee JJ, Lotvall JO. (2007) Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol*; 9: 654-659.

Van Damme EJM, Lannoo N, Peumans WJ. (2008) Plant lectins. *Advances in Botanical Research Incorporation Advances in Plant Pathology* 48:107–209. DOI 10.1016/S0065-2296(08)00403-5.

Vandenberg LN, Chahoud I, Heindel JJ, Padmanabhan V, Paumgarten FJ, Schoenfelder G. (2010) Urinary, circulating and tissue biomonitoring studies indicate widespread exposure to bisphenol A. *Environ Health Perspect*, 118:1055–1070. doi: 10.1289/ehp.0901716.

Vandenberg LN, Maffini MV, Sonnenschein C, Rubin BS, Soto AM. (2009). *Endocr Rev*, 30:75–95. doi: 10.1210/er.2008-0021.

Varki A, Cummings RD, Esko JD, Stanley P, Hart GW, Aebi M, Darvill AG, Kinoshita T, Packer NH, Prestegard JH, Schnaar RL, Seeberger PH. (eds) (2009) *Essentials of Glycobiology* 2nd edn (Cold Spring Harbor Laboratory Press).

Vermeersch KA, Styczynski MP. (2013) Applications of metabolomics in cancer research. *J Carcinog*; 12: 9.

Wakatsuki T, Irisawa A, Bhutani MS, Hikichi T, Shibukawa G, Takagi T, Yamamoto G, Takahashi Y, Yamada Y, Watabane K, Obara K, Suzuki T, Sato T. (2005) Comparative study of diagnostic value of cytologic sampling by endoscopic ultrasonography-guided fine-needle aspiration and that by endoscopic retrograde pancreatography for the management of pancreatic mass without biliary stricture. *J Gastroenterol Hepatol*;20:1707-11. <https://doi.org/10.1111/j.1440-1746.2005.03900.x>

Wang M, Long RE, Comunale MA, Junaidi O, Marrero J, Di Bisceglie AM, Block TM, Mehta AS. (2009) Novel fucosylated biomarkers for the early detection of hepatocellular carcinoma. *Cancer Epidemiol Biomarkers Prev*; 18: 1914-1921. PMID: 19454616 DOI: 10.1158/1055-9965.EPI-08-0980

Wang, Yingxue, Li, Zhihua, Zheng, Shangyou, Zhou, Yu, Zhao, Lei, Ye, Huilin, Zhao, Xiaohui, Gao, Wenchao, Fu, Zhiqiang, Zhou, Quanbo, Liu, Yimin, Chen, Rufu. (2015). Expression profile of long non-coding RNAs in pancreatic cancer and their clinical significance as biomarkers. *Oncotarget*. 6. 10.18632/oncotarget.5533.

Warburg, O. (1956) On the origin of cancer cells. *Science* 123, 309–314.

Wingren C, Sandstrom A, Segersvard R, Carlsson A, Andersson R, Lohr M, Borrebaeck CA. (2012) Identification of serum biomarker signatures associated with pancreatic cancer. *Cancer Res*; 72: 2481-2490.

Wong D, Ko AH, Hwang J, Venook AP, Bergsland Ek, Tempero MA. (2008) Serum CA19-9 decline compared to radiographic response as a surrogate for clinical outcomes in patients with metastatic pancreatic cancer receiving chemotherapy. *Pancreas*; 37:269-74. doi: 10.1097/MPA.0b013e31816d8185.

World Cancer Report, International Agency for Research on Cancer, 2014

World cancer research fund international/ American institute for cancer research (2018) Diet, Nutrition, Physical Activity and Cancer: A Global Perspective. Continuous Update Project Expert Report 2018. Available at [dietandcancerreport.org](http://dietandcancerreport.org)

World Cancer Research Fund/American Institute for Cancer Research (2018) Diet, Nutrition, Physical Activity and Cancer: a Global Perspective. Continuous Update Project Expert Report 2018. Available at [dietandcancerreport.org](http://dietandcancerreport.org)

World Health Organization website, <https://www.who.int/topics/epidemiology/en/> (accessed 2 February 2, 2019)

Xu J, Cao Z, Liu W, You L, Zhou L, Wang C, Lou W, Sun B, Miao Y, Liu X, Zhang T, Zhao Y. (2016) Plasma miRNAs effectively distinguish patients with pancreatic cancer from controls: a Multicenter Study. *Ann Surg*; 263: 1173- 1179.

Xue A, Scarlett CJ, Chung L, Butturini G, Scarpa A, Gandy R, Wilson SR, Baxter RC, Smith RC (2010) Discovery of serum biomarkers for pancreatic adenocarcinoma using proteomic analysis. *Br J Cancer* 103: 391 – 400.

Xue Y, Abou Tayoun AN, Abo KM, Pipas JM, Gordon SR, Gardner TB, Barth RJ Jr, Suriawinata AA, Tsongalis GJ. (2013) MicroRNAs as diagnostic markers for pancreatic ductal adenocarcinoma and its precursor, pancreatic intraepithelial neoplasm. *Cancer Genet*; 206: 217- 221.

Yachida S, Jones S, Bozic I, Antal T, Leary R, Fu B, Kamiyama M, Hruban RH, Eshleman JR, Nowak MA, Velculescu VE, Kinzler KW, Vogelstein B, Iacobuzio-Donahue CA. (2010) Distant metastasis occurs late during the genetic evolution of pancreatic cancer. *Nature*; 467: 1114-1117.

Yanagisawa A, Ohtake K, Ohashi K, Hori M, Kitagawa T, Sugano H, Kato Y. (1993) Frequent c-Ki-ras oncogene activation in mucous cell hyperplasias of pancreas suffering from chronic inflammation. *Cancer Res*; 53: 953-956.

Yang F, Jin C, Subedi S, Lee CL, Wang Q, Jiang Y, Li J, Di Y, Fu D. Emerging inorganic nanomaterials for pancreatic cancer diagnosis and treatment. *Cancer Treat Rev* 2012; 38: 566-579.

Yang S, Che SP, Kurywchak P, Tavormina JL, Gansmo LB, Correa de Sampaio P, Tachezy M, Bockhorn M, Gebauer F, Haltom AR, Melo SA, LeBleu VS, Kalluri R. (2017) Detection of mutant KRAS and TP53 DNA in circulating exosomes from healthy individuals and patients with pan- creatic cancer. *Cancer Biol Ther*; 18: 158-165.

Yi JM, Guzzetta AA, Bailey VJ, Downing SR, Van Neste L, Chiappinelli KB, Keeley BP, Stark A, Herrera A, Wolfgang C, Pappou EP, Iacobuzio- Donahue CA, Goggins MG, Herman JG, Wang TH, Baylin SB, Ahuja N. (2013) Novel methylation biomarker panel for the early detection of pancreatic cancer. *Clin Cancer Res*; 19: 6544-6555.

Yoshinaga T, Niou T, Niihara T, Kajiya Y, Hori E, Tomiyoshi A, Tokudome E, Nishimata H, Takei T, Yoshida M. (2018) Angiopoietin-like Protein 2 is a Useful Biomarker for Pancreatic Cancer that is Associated with Type 2 Diabetes Mellitus and Inflammation. *J Cancer*. 2018 Nov 25;9(24):4736-4741.

Young M, Wagner P, Ghosh S, Rinaudo J, Baker S, Zaret K, Goggins M, Srivastava S. (2018) Validation of Biomarkers for Early Detection of Pancreatic Cancer: Summary of the Alliance of Pancreatic Cancer Consortia for Biomarkers for Early Detection Workshop, Pancreas. 2018 February; 47(2): 135–141.

Yu KH, Rustgi AK, Blair IA. (2005) Characterization of proteins in human pancreatic cancer serum using differential gel electrophoresis and tandem mass spectrometry. *J Proteome Res* 4: 1742 – 1751.

Yuan C, Clish CB, Wu C, Mayers JR, Kraft P, Townsend MK, Zhang M, Tworoger SS, Bao Y, Qian ZR, Robinson DA, Ng K, Giovannucci EL, Ogino S, Stampfer MJ, Gaziano JM, Ma J, Ses- so HD, Anderson GL, Cochrane BB, Manson JE, Torrence ME, Kimmelman AC, Amundadottir LT, Vander Heiden MG, Fuchs CS, Wolpin BM. (2016) Circulating metabolites and survival among patients with pancreatic cancer. *J Natl Cancer Inst*; 108: djv409. doi: 10.1093/jnci/djv409.

Yue T, Goldstein IJ, Hollingsworth MA, Kaul K, Brand RE, Haab BB. (2009) The prevalence and nature of glycan alterations on specific proteins in pancreatic cancer patients revealed using antibody-lectin sandwich arrays. *Mol Cell Proteomics*; 8: 1697-1707. PMID: 19377061 DOI: 10.1074/mcp.M900135-MCP200

Yue T, Partyka K, Maupin KA, Hurley M, Andrews P, Kaul K, Moser AJ, Zeh H, Brand RE, Haab BB. (2011) Identification of blood-protein carriers of the CA 19-9 antigen and characterization of prevalence in pancreatic diseases. *Proteomics*, 11, 3665–3674.

Zeh HJ, Winikoff S, Landsittel DP, Gorelik E, Marrangoni AM, Velikokhatnaya L, Winans MT, Lee K, Moser A, Bartlett D, Lotze MT, Siegfried JM, Whitcomb D,



Papacristou G, Slivka A, Bigbee WL, Lokshin AE. (2005) Multianalyte profiling of serum cytokines for detection of pancreatic cancer. *Cancer Biomark*; 1: 259-269.

Zhang G, He P, Tan H, Budhu A, Gaedcke J, Ghadimi BM, Ried T, Yfantis HG, Lee DH, Maitra A, Hanna N, Alexander HR, Hussain SP. (2013) Integration of metabolomics and transcriptomics revealed a fatty acid network exerting growth inhibitory effects in human pancreatic cancer. *Clin Cancer Res*; 19: 4983- 4993.

Zhang Y, Yang J, Li H, Wu Y, Zhang H, Chen W. (2015) Tumor markers CA19-9, CA242 and CEA in the diagnosis of pancreatic cancer: a meta- analysis. *Int J Clin Exp Med*; 8: 11683- 11691.

Zhang L, Sanagapalli S, Stoita A. (2018a) Challenges in diagnosis of pancreatic cancer, *World J Gastroenterol*; 24(19): 2047-2060

Zhang X, Shi S, Zhang B, Ni Q, Yu X, Xu J. (2018b) Circulating biomarkers for early diagnosis of pancreatic cancer: facts and hopes, *Am J Cancer Res*; 8(3):332-353

Zhao J, Patwa TH, Qiu W, Shedden K, Hinderer R, Misek DE, Anderson MA, Simeone DM, Lubman DM. (2007) Glycoprotein microarrays with multi-lectin detection: unique lectin binding patterns as a tool for classifying normal, chronic pancreatitis and pancreatic cancer sera. *J Proteome Res*; 6: 1864-1874 [PMID: 17428079 DOI: 10.1021/pr070062p]

Zhao J, Qiu W, Simeone DM, Lubman DM. (2007) N-linked glycosylation profiling of pancreatic cancer serum using capillary liquid phase separation coupled with mass spectrometric analysis. *J Proteome Res*; 6: 1126-1138. PMID: 17249709 DOI: 10.1021/pr0604458

Zhao Y, Sato Y, Isaji T, Fukuda T, Matsumoto A, Miyoshii E, Gu J, Taniguchi N. (2008) Branched N-glycans regulate the biological functions of integrins and cadherins. *FEBS J*. 275, 1939–1948. doi: 10.1111/j.1742-4658.2008.06346.x.


Zheng L, Xue J, Jaffee EM, Habtezion A. (2013) Role of immune cells and immune-based therapies in pancreatitis and pancreatic ductal adenocarcinoma. *Gastroenterology*, 144:1230–1240.

Zhou W, Capello M, Fredolini C, Racanicchi L, Piemonti L, Liotta LA, Novelli F, Petricoin EF. (2012) Proteomic analysis reveals Warburg effect and anomalous metabolism of glutamine in pancreatic cancer cells. *J Proteome Res*; 11: 554-563.


















Zoller M. (2013) Pancreatic cancer diagnosis by free and exosomal miRNA. *World J Gastrointest Pathophysiol*; 4: 74-90.


# Annexes

## A1. The evidence for cancer risk: summary matrix



Part 1. Adapted from World Research Fund, 2018

Conclusions Key		Exposure Group Key	
	Convincing decreases risk		Convincing increases risk
	Probable decreases risk		Probable increases risk
	Limited – suggestive decreases risk		Limited – suggestive increases risk
			Substantial effect on risk unlikely
	Wholegrains, vegetables and fruit		Other dietary exposures
	Meat, fish and dairy products		Physical activity
	Preservation and processing of foods		Body fatness and weight gain
	Non-alcoholic drinks		Height and birthweight
	Alcoholic drinks		Lactation/having been breastfed



Part 2. Adapted from World Cancer Research Fund, 2018

Conclusions Key				Exposure Group Key			
<div></div>	Convincing decreases risk	<div></div>	Convincing increases risk	<div></div>	Wholegrains, vegetables and fruit	<div></div>	Other dietary exposures
<div></div>	Probable decreases risk	<div></div>	Probable increases risk	<div></div>	Meat, fish and dairy products	<div></div>	Physical activity
<div></div>	Limited – suggestive decreases risk	<div></div>	Limited – suggestive increases risk	<div></div>	Preservation and processing of foods	<div></div>	Body fatness and weight gain
		<div></div>	Substantial effect on risk unlikely	<div></div>	Non-alcoholic drinks	<div></div>	Height and birthweight
				<div></div>	Alcoholic drinks	<div></div>	Lactation/having been breastfed